

**AN ALMOST SURE INVARIANCE PRINCIPLE  
FOR RENORMALIZED INTERSECTION LOCAL TIMES**

**Richard F. Bass**<sup>1</sup> and **Jay Rosen**<sup>2</sup>

**Abstract.** Let  $\tilde{\beta}_k(n)$  be the number of self-intersections of order  $k$ , appropriately renormalized, for a mean zero random walk  $X_n$  in  $\mathbb{Z}^2$  with  $2 + \delta$  moments. On a suitable probability space we can construct  $X_n$  and a planar Brownian motion  $W_t$  such that for each  $k \geq 2$

$$|\tilde{\beta}_k(n) - \tilde{\gamma}_k(n)| = O(n^{-a}), \quad \text{a.s.}$$

for some  $a > 0$  where  $\tilde{\gamma}_k(n)$  is the renormalized self-intersection local time of order  $k$  at time 1 for the Brownian motion  $W_{nt}/\sqrt{n}$ .

**Keywords and Phrases:** intersection local time, invariance principle, random walks, strong invariance, self-intersection local time

**AMS 2000 Subject Classifications:** Primary 60F17; Secondary 60J55

Submitted to EJP on July 8, 2004. Final version accepted on January 28, 2005.

Short title: Self-intersection local time

---

<sup>1</sup> Research partially supported by NSF grant DMS-0244737.

<sup>2</sup> Research partially supported by grants from the NSF and from PSC-CUNY.

## 1. Introduction.

If  $\{W_t; t \geq 0\}$  is a planar Brownian motion with density  $p_t(x)$ , for  $x \in \mathbb{R}^2$  set  $\gamma_{1,\epsilon}(t, x) = t$  and for  $k \geq 2$  and  $x = (x_2, \dots, x_k) \in (\mathbb{R}^2)^{k-1}$  let

$$\gamma_{k,\epsilon}(t, x) = \int_{0 \leq t_1 \leq \dots \leq t_k < t} \prod_{i=2}^k p_\epsilon(W_{t_i} - W_{t_{i-1}} - x_i) dt_1 \cdots dt_k.$$

When  $x_i \neq 0$  for all  $i$  the limit

$$\gamma_k(t, x) = \lim_{\epsilon \rightarrow 0} \gamma_{k,\epsilon}(t, x)$$

exists and for any bounded continuous function  $F(x)$  on  $\mathbb{R}^{2(k-1)}$  we have

$$\int F(x) \gamma_k(t, x) dx = \int_{0 \leq t_1 \leq \dots \leq t_k < t} F(W_{t_2} - W_{t_1}, \dots, W_{t_k} - W_{t_{k-1}}) dt_1 \cdots dt_k. \quad (1.1)$$

(Here we may arbitrarily specify that  $\gamma_k(t, x) = \infty$  if any  $x_i = 0$ .) When  $x_i \neq 0$  for all  $i$  define the renormalized intersection local times as

$$\tilde{\gamma}_k(t, x) = \sum_{A \subseteq \{2, \dots, k\}} (-1)^{|A|} \left( \prod_{i \in A} \frac{1}{\pi} \log(1/|x_i|) \right) \gamma_{k-|A|}(t, x_{A^c})$$

where  $x_{A^c} = (x_{i_1}, \dots, x_{i_{k-|A|}})$  with  $i_1 < i_2 < \dots < i_{k-|A|}$  and  $i_j \in \{2, \dots, k\} - A$  for each  $j$ , that is, the vector  $(x_2, \dots, x_k)$  with all terms that have indices in  $A$  deleted. (When  $A = \{2, \dots, k\}$  so that  $k - |A| = 1$  and  $A^c = \emptyset$  we set  $\gamma_{k-|A|}(t, x_{A^c}) = \gamma_{1,\epsilon}(t) = t$ .) It is known that the  $\tilde{\gamma}_k(t, x)$  have a continuous extension to all  $\mathbb{R}_+^1 \times \mathbb{R}^{k-1}$ ; see [3].

Renormalized self-intersection local time was originally studied by Varadhan [20] for its role in quantum field theory. In Rosen [18] it is shown that  $\tilde{\gamma}_k(t, 0)$  can be characterized as the continuous process of zero quadratic variation in the decomposition of a natural Dirichlet process. Renormalized intersection local time turns out to be the right tool for the solution of certain ‘‘classical’’ problems such as the asymptotic expansion of the area of the Wiener sausage in the plane and the range of random walks, [5], [9], [10]. For further work on renormalized self-intersection local times see Dynkin [7], Le Gall [11], Bass and Khoshnevisan [3], Rosen [17] and Marcus and Rosen [14].

Let  $\xi_i$  be i.i.d. random variables with values in  $\mathbb{Z}^2$  that are mean 0, with covariance matrix equal to the identity, and with  $2 + \delta$  moments. Let us suppose the  $\xi_i$  are symmetric and are strongly aperiodic. Let  $X_n$  be the random walk, that is,  $X_n = \sum_{i=1}^n \xi_i$ . Let  $p(n, x, y)$  be the transition probabilities. Let  $B_1(n, x) = n$  and for  $x \in \mathbb{Z}^2$  set

$$B_2(n, x) = \sum_{0 \leq i_1 < i_2 \leq n} 1_{(X_{i_2} = X_{i_1} + x)}.$$

More generally, for  $x = (x_2, \dots, x_k) \in (\mathbb{Z}^2)^{k-1}$  let

$$B_k(n, x) = \sum_{0 \leq i_1 < i_2 < \dots < i_k \leq n} \prod_{j=2}^k 1_{(X_{i_j} = X_{i_{j-1}} + x_j)}.$$

Note that  $B_k(n, x) = 0$  for all  $n < k - 1$ .

With  $e_1 = (1, 0)$ , let

$$G(x) = \sum_{n=1}^{\infty} [p(n, 0, x) - p(n, 0, e_1)],$$

and set  $G_n(x) = G(x) - G(\sqrt{n}e_1)$ . Let

$$\tilde{B}_k(n, x) = \sum_{A \subset \{2, \dots, k\}} (-1)^{|A|} \left( \prod_{i \in A} G_n(x_i) \right) B_{k-|A|}(n, x_{A^c}). \quad (1.2)$$

In particular we have

$$\tilde{B}_2(n, x) = B_2(n, x) - G_n(x)n. \quad (1.3)$$

Finally we define the renormalized intersection local times for our random walk by

$$\tilde{\beta}_k(n, x) = \frac{1}{n} \tilde{B}_k(n, x\sqrt{n}). \quad (1.4)$$

In particular we have

$$\tilde{\beta}_2(n, x) = \frac{1}{n} B_2(n, x\sqrt{n}) - G_n(x\sqrt{n}). \quad (1.5)$$

We note from P12.3 of [19] that for  $x \neq 0$

$$\lim_{n \rightarrow \infty} G_n(x\sqrt{n}) = \lim_{n \rightarrow \infty} G(x\sqrt{n}) - G(\sqrt{n}e_1) = \frac{1}{\pi} \log(1/|x|). \quad (1.6)$$

We know we can find a version of our random walk and a Brownian motion  $W_t$  such that

$$\sup_{s \leq 1} |X_s^n - W_s^n| = o(n^{-\zeta}), \quad a.s. \quad (1.7)$$

for some  $\zeta > 0$  where  $X_t^n = X_{[nt]}/\sqrt{n}$  and  $W_t^n = W_{nt}/\sqrt{n}$ ; see, [6], Theorem 3, for example. Let  $\gamma_k(1, x, n)$  and  $\tilde{\gamma}_k(1, x, n)$  be the intersection local times and renormalized intersection local times up to time 1 of order  $k$ , resp., for the Brownian motion  $W_t^n$ . In this paper we prove the following theorem.

**Theorem 1.1.** *Let  $X_n = \xi_1 + \cdots + \xi_n$  be a random walk in  $\mathbb{Z}^2$ , where the  $\xi_i$  are i.i.d., mean 0, with covariance matrix equal to the identity, with  $2 + \delta$  moments for some  $\delta > 0$ , symmetric, and strongly aperiodic. On a suitable probability space we can construct  $\{X_n; n \geq 1\}$  and a planar Brownian motion  $\{W_t; t \geq 0\}$  and we can find  $\eta > 0$  such that for each  $k \geq 2$*

$$|\tilde{\beta}_k(n, 0) - \tilde{\gamma}_k(1, 0, n)| = o(n^{-\eta}), \quad a.s.$$

For related work see [4], [5], [16]. In fact, [5] provided much of the motivation for this work; in that paper we proved a strong invariance principle with respect to the  $L^2$  norm.

We give a brief overview of the proof. There is an equation similar to (1.1) when  $\gamma_k$  is replaced by  $\tilde{\gamma}_k$ , and also when it is replaced by  $\tilde{\beta}_k$ . Since by (1.7) we have  $X_s^n$  close to  $W_s^n$  for  $n$  large, we are able to conclude that  $\int F(x)\tilde{\gamma}_k(1, x) dx$  is close to  $\int F(x)\tilde{\beta}_k(n, x) dx$  for  $n$  large. If  $F$  is smooth, with integral 1 and support in a small neighborhood of the origin, then by the continuity of  $\tilde{\gamma}_k(t, x)$  in  $x$ , which is proved in [3], we see that  $\int F(x)\tilde{\gamma}_k(1, x) dx$  is not far from  $\tilde{\gamma}_k(1, 0)$ . If we had a similar result for  $\tilde{\beta}_k$ , we would then have that  $\int F(x)\tilde{\beta}_k(n, x) dx$  is not far from  $\tilde{\beta}_k(n, 0)$ , and we would have our proof. So our strategy is to obtain good estimates on  $|\tilde{\beta}_k(n, x) - \tilde{\beta}_k(n, 0)|$ . Because of the rate of convergence in (1.7), it turns out we are able to avoid having to find the sharpest estimates on the difference, which simplifies the proof considerably.

Our main tool in obtaining the desired estimates is Proposition 3.2. This proposition may be of independent interest. It has been known for a long time that one way of proving  $L^p$  estimates for a continuous increasing process is to prove corresponding estimates for the potential. It is not as well known that one can do the same for continuous processes of bounded variation provided one has some control on the total variation; see, e.g., [1] or [3]. Proposition 3.2 is the discrete time analogue of this result, and is proved in a similar way. Unlike the continuous time version, here it is also necessary to have control on the differences of successive terms.

Section 2 has some estimates on the potential kernel for random walks in the plane, while Section 3 has the proof of the stochastic calculus results we need. Theorem 1.1 in the case when  $k = 2$  is proved in Section 4, with the proofs of some lemmas postponed to Section 5. We treat the case  $k = 2$  separately for simplicity of exposition. The description of the potentials of intersection local times of random walks in the  $k > 2$  case is a bit different than in the  $k = 2$  case and this is described in Section 6. Theorem 1.1 in the  $k > 2$  case is proved in Section 7, with the proofs of some lemmas given in Sections 8 and 9. Finally in Section 10 we give an extension of our results to  $L^2$  convergence, and more importantly, make a correction to the proof of one of the propositions in [3]. An Appendix contains the detailed proof of that correction. Throughout this paper we use the letter  $c$  to denote finite positive constants whose exact value is unimportant and which may vary

from line to line.

## 2. Estimates for random walks.

In this section we prove some estimates for the potential kernel of a random variable. See the forthcoming book by Lawler [13] for further information. Let  $\bar{G}$  be the potential kernel for  $X$ . Recall that in 2 dimensions, since  $X$  is recurrent, the potential kernel is defined somewhat differently than in higher dimensions, and is defined by

$$\bar{G}(x) = \sum_{n=0}^{\infty} [p(n, 0, x) - p(n, 0, e_1)],$$

where  $e_1 = (1, 0)$ . (Note  $e_1$  can be replaced by any fixed point.) For us it will be more convenient to work with

$$G(x) = \sum_{n=1}^{\infty} [p(n, 0, x) - p(n, 0, e_1)],$$

which, since  $p(0, 0, 0) = 1$  and  $p(0, 0, e_1) = 0$ , differs from  $\bar{G}(x)$  by  $1_{\{0\}}(x)$ . By Spitzer [19], p. 75, we have

$$p(n, 0, x) \leq c/n. \tag{2.1}$$

By [4], Proposition 2.1, if the  $\xi_i$  are strongly aperiodic, then

$$|p(n, 0, x) - p(n, 0, y)| \leq \frac{c|x - y|}{n^{3/2}}. \tag{2.2}$$

**Proposition 2.1.** *Suppose the  $\xi_i$  have  $2 + \delta$  moments. Then  $G(x)$  exists and  $|G(x)| \leq c(1 + \log^+ |x|)$ .*

**Proof.** Using (2.2), we have that

$$|G(0)| \leq \sum_{n=1}^{\infty} \frac{c}{n^{3/2}}$$

is finite. The rest of the assertions follow from

$$\begin{aligned} |G(x)| &\leq \sum_{n=1}^{|x|^2} \frac{c}{n} + \sum_{n=|x|^2+1}^{\infty} \frac{c|x - e_1|}{n^{3/2}} \\ &\leq c + c \log |x| + c \frac{|x|}{(|x|^2 + 1)^{1/2}}. \end{aligned}$$

□

**Proposition 2.2.** For some  $c < \infty$

$$|G(x) - G(y)| \leq c \left( \frac{|x - y|}{(1 + |x|) \wedge (1 + |y|)} \right)^{2/3}, \quad x, y \in \mathbb{Z}^2.$$

**Proof.** By [19], P7.10,

$$p(j, 0, x) \leq \frac{c}{|x|^2}. \quad (2.3)$$

Since  $p(j, 0, 0) \leq 1$ , then we have

$$p(j, 0, x) \leq \frac{c}{1 + |x|^2}. \quad (2.4)$$

Suppose  $0 < |x| \leq |y|$ . Let us set  $R$  in a moment. Using (2.4) for  $j \leq R$  and (2.2) for  $j > R$ , we have that

$$|G(x) - G(y)| \leq \sum_{j=1}^R \frac{c}{1 + |x|^2} + \sum_{j=R+1}^{\infty} \frac{c|x - y|}{j^{3/2}} \leq \frac{cR}{1 + |x|^2} + \frac{c|x - y|}{R^{1/2}}.$$

If we select  $R$  so that

$$\frac{R}{1 + |x|^2} = \frac{|x - y|}{R^{1/2}}, \quad \text{i.e.,} \quad R^{3/2} = (1 + |x|^2)(|x - y|),$$

the result follows. Since  $G(0)$  is finite and  $|G(x)| \leq c \log(1 + |x|) \leq |x|^{2/3}$ , the result holds when either  $x$  or  $y$  is 0, as well.  $\square$

**Lemma 2.3.** For some constant  $\kappa$  and any  $\rho < \delta/2$ ,

$$G(x) = \kappa + \frac{1}{\pi} \log(1/|x|) + O(|x|^{-\rho}), \quad x \in \mathbb{Z}^2.$$

**Proof.** Let us begin with the proof of Proposition 3.1 in [2]. We have for  $\delta > 0$

$$\left| e^{i\alpha \cdot x / \sqrt{n}} - 1 - \frac{\alpha \cdot x}{\sqrt{n}} - \frac{|\alpha \cdot x|^2}{2n} \right| \leq c \left| \frac{\alpha \cdot x}{\sqrt{n}} \right|^{2+\delta}.$$

So if  $\phi$  is the characteristic function of a random vector with finite  $2 + \delta$  moments, mean 0, and the identity as its covariance matrix, then

$$\phi(\alpha/\sqrt{n}) = 1 - \frac{|\alpha|^2}{2n} + E_1(\alpha, n),$$

with

$$|E_1(\alpha, n)| \leq c(|\alpha|/\sqrt{n})^{2+\delta}. \quad (2.5)$$

Applying this also for the characteristic function of a standard normal vector,

$$e^{-|\alpha|^2/2n} = 1 - \frac{|\alpha|^2}{2n} + E_2(\alpha, n),$$

where  $E_2(\alpha, n)$  has the same bound as  $E_1(\alpha, n)$ . If we use this in place of the display in the middle of page 473 of [2], we obtain

$$I_1^{(n)} \leq cn^{-\delta/2}(\log n)^{(4+\delta)/2}.$$

So if  $E(n, x) = |p(n, 0, x) - (2\pi n)^{-1}e^{-|x|^2/2n}|$ , following the proof in [2] we obtain

$$\sup_x E(n, x) \leq cn^{-1-(\delta/2)}(\log n)^{(4+\delta)/2}.$$

Let us choose  $\delta' < \delta$ . We then have

$$\sup_x E(n, x) \leq cn^{-1-(\delta'/2)}. \quad (2.6)$$

Recall  $G(x) = \sum_{k=1}^{\infty} [p(k, 0, x) - p(k, 0, e_1)]$ . It is shown in the proof of Theorem 1.6.2 in [12] that for some constant  $\kappa$

$$\sum_{k=1}^{\infty} [q(k, 0, x) - q(k, 0, e_1)] = \kappa + \frac{1}{\pi} \log(1/|x|) + O(|x|^{-1}),$$

where  $q(k, x, y) = (2\pi k)^{-1}e^{-|x-y|^2/2k}$ . Thus, to prove the lemma, it suffices to prove

$$\sum_{k=1}^{\infty} |p(k, x, 0) - q(k, x, 0)| = O(|x|^{-\rho}) \quad (2.7)$$

for any  $\rho < \delta/2$ .

To establish (2.7), use [15], p. 60 to observe that

$$p(k, x, 0) \leq \mathbb{P}(|X_k| > |x|) \leq \frac{\mathbb{E}|X_k|^{2+\delta}}{|x|^{2+\delta}} \leq c \frac{k^{1+\delta/2}}{|x|^{2+\delta}}$$

and a similar estimate is easily seen to hold for  $q(k, x, 0)$ . Therefore, using (2.6) and setting  $R = |x|$ ,

$$\begin{aligned} \sum_{k=1}^{\infty} |p(k, x, 0) - q(k, x, 0)| &\leq \sum_{k=1}^R [p(k, x, 0) + q(k, x, 0)] \\ &\quad + \sum_{k=R}^{\infty} |p(k, x, 0) - q(k, x, 0)| \\ &\leq \sum_{k=1}^R c \frac{k^{1+\delta/2}}{|x|^{2+\delta}} + \sum_{k=R}^{\infty} c \frac{1}{k^{1+\delta'/2}} \\ &\leq c \frac{R^{2+\delta/2}}{|x|^{2+\delta}} + c \frac{1}{R^{\delta'/2}} \\ &\leq c|x|^{-\rho}. \end{aligned}$$

□

### 3. Stochastic calculus.

We will use the following propositions; these may be of independent interest. Propositions 3.1 and 3.2 and their proofs are the discrete time analogues of Propositions 6.1 and 6.2 of [3].

**Proposition 3.1.** *Let  $A_n$  be an adapted increasing sequence of random variables with  $A_0 = 0$  and  $A_\infty = \sup_n A_n$  finite. Suppose that*

$$Y = \sup_n (A_n - A_{n-1})$$

and  $W$  is a random variable such that

$$\mathbb{E}[A_\infty - A_n \mid \mathcal{F}_n] \leq \mathbb{E}[W \mid \mathcal{F}_n]$$

for all  $n$ . Then for each integer  $p$  larger than 1 there exists a constant  $c$  such that

$$\mathbb{E} A_\infty^p \leq cp^p (\|W\|_p + \|Y\|_p)^p.$$

**Proof.** Since  $A_n$  is increasing,

$$pA_n^{p-1} \geq A_n^{p-1} + A_n^{p-2}A_{n-1} + \cdots + A_nA_{n-1}^{p-2} + A_{n-1}^{p-1}.$$

Multiplying by  $A_n - A_{n-1}$ , we obtain

$$(A_n - A_{n-1})pA_n^{p-1} \geq A_n^p - A_{n-1}^p.$$

Summing over  $n$  we obtain

$$p \sum_{n=1}^{\infty} (A_n - A_{n-1}) A_n^{p-1} \geq A_\infty^p. \quad (3.1)$$

On the other hand, applying the general summation formula

$$A_\infty B_\infty = \sum_{n=1}^{\infty} A_n (B_n - B_{n-1}) + \sum_{n=1}^{\infty} (A_n - A_{n-1}) B_{n-1}$$

with  $B_n = A_{n+1}^{p-1}$  we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} (A_n - A_{n-1}) A_n^{p-1} &= A_\infty^p - \sum_{n=1}^{\infty} A_n (A_{n+1}^{p-1} - A_n^{p-1}) \\ &= \sum_{n=1}^{\infty} (A_\infty - A_n) (A_{n+1}^{p-1} - A_n^{p-1}) + A_\infty A_1^{p-1}. \end{aligned} \quad (3.2)$$



Here we used the fact that

$$\sum_{n=1}^{\infty} A_{\infty}(A_{n+1}^{p-1} - A_n^{p-1}) = A_{\infty}^p - A_{\infty}A_1^{p-1}.$$

Combining (3.1) and (3.2) we obtain

$$A_{\infty}^p \leq p \sum_{n=1}^{\infty} (A_{\infty} - A_n)(A_{n+1}^{p-1} - A_n^{p-1}) + pA_{\infty}A_1^{p-1}. \quad (3.3)$$

Now suppose for the moment that  $Y$  is bounded and  $A_n = A_{n_0}$  for all  $n \geq n_0$  for some  $n_0$ . We have

$$\begin{aligned} & \sum_{n=1}^{\infty} (A_{\infty} - A_n)(A_{n+1}^{p-1} - A_n^{p-1}) \\ &= \sum_{n=1}^{\infty} (A_{\infty} - A_{n+1})(A_{n+1}^{p-1} - A_n^{p-1}) + \sum_{n=1}^{\infty} (A_{n+1} - A_n)(A_{n+1}^{p-1} - A_n^{p-1}). \end{aligned} \quad (3.4)$$

and

$$\sum_{n=1}^{\infty} (A_{n+1} - A_n)(A_{n+1}^{p-1} - A_n^{p-1}) \leq YA_{\infty}^{p-1}.$$

But,  $A_1 \leq A_{\infty}$  and also  $A_1 \leq Y$  so that

$$A_{\infty}A_1^{p-1} \leq YA_{\infty}^{p-1}.$$

We write

$$\begin{aligned} \mathbb{E} \sum_{n=1}^{\infty} (A_{\infty} - A_{n+1})(A_{n+1}^{p-1} - A_n^{p-1}) &= \mathbb{E} \sum_{n=1}^{\infty} \mathbb{E}[A_{\infty} - A_{n+1} \mid \mathcal{F}_{n+1}](A_{n+1}^{p-1} - A_n^{p-1}) \\ &\leq \mathbb{E} \sum_{n=1}^{\infty} \mathbb{E}[W \mid \mathcal{F}_{n+1}](A_{n+1}^{p-1} - A_n^{p-1}) \\ &= \mathbb{E} \sum_{n=1}^{\infty} W(A_{n+1}^{p-1} - A_n^{p-1}) \\ &\leq \mathbb{E}[WA_{\infty}^{p-1}]. \end{aligned}$$

Therefore using Hölder's inequality,

$$\mathbb{E} A_{\infty}^p \leq p(\|W\|_p + 2\|Y\|_p)(\mathbb{E} A_{\infty}^p)^{1-\frac{1}{p}}.$$

Our temporary assumptions on  $A$  allow us to divide both sides by  $(\mathbb{E} A_{\infty}^p)^{1-\frac{1}{p}}$  to obtain our result in this special case.

In general, look at

$$A'_n = \sum_{j=1}^n ((A_j - A_{j-1}) \wedge K)$$

and apply the above to  $A''_n = A'_{n \wedge n_0}$ ; note that  $A''$  will satisfy the hypotheses with the same  $W$  and  $Y$ . Then let  $K \uparrow \infty$  and next  $n_0 \uparrow \infty$  and use monotone convergence.  $\square$

**Proposition 3.2.** *Suppose  $Q_n^1$  and  $Q_n^2$  are two adapted nonnegative increasing sequences. Suppose  $Q_n = Q_n^1 - Q_n^2$ ,  $H_n + Q_n$  is a martingale that is 0 at time 0,*

$$Z = \sup_n |H_n|,$$

$$Y = \sup_n [(Q_n^1 - Q_{n-1}^1) + (Q_n^2 - Q_{n-1}^2)],$$

and

$$W = Q_\infty^1 + Q_\infty^2.$$

Then there exists  $c$  such that for  $p > 1$

$$\begin{aligned} \mathbb{E} \sup_n |Q_n|^{2p} &\leq c^p p^{4p} \left[ \mathbb{E} Z^{2p} + (\mathbb{E} Z^{2p})^{1/2} (\mathbb{E} W^{2p})^{1/2} \right. \\ &\quad \left. + (\mathbb{E} Y^{2p})^{1/2} (\mathbb{E} W^{2p})^{1/2} \right]. \end{aligned} \tag{3.5}$$

**Proof.** There is nothing to prove unless  $\mathbb{E} W^{2p} < \infty$ . Since  $\sup_n Q_n \leq W$ , all the random variables that follow will satisfy the appropriate integrability conditions. Let us temporarily assume that there exists  $n_0$  such that  $Q_n^i = Q_{n_0}^i$  if  $n \geq n_0, i = 1, 2$ .

Let

$$V_m = \mathbb{E}[Q_\infty - Q_m | \mathcal{F}_m], \quad M_m = \mathbb{E}[Q_\infty | \mathcal{F}_m].$$

Note that  $V_\infty = 0$ ,  $M_m$  is a martingale, and  $Q_m = M_m - V_m$ . In fact, in view of our temporary assumption,  $V_m = 0$  if  $m \geq n_0$ .

Our first observation is that since

$$V_m = \mathbb{E}[Q_\infty - Q_m | \mathcal{F}_m] = \mathbb{E}[H_m - H_\infty | \mathcal{F}_m],$$

then

$$|V_m| \leq 2\mathbb{E}[Z | \mathcal{F}_m]. \tag{3.6}$$

By Doob's inequality,

$$\mathbb{E} \sup_n V_n^p \leq c\mathbb{E} Z^p. \tag{3.7}$$

We will use

**Lemma 3.3.**

$$\mathbb{E} \left( \sum_{n=0}^{\infty} (M_{n+1} - M_n)^2 \right)^p \leq c^p p^{2p} (\mathbb{E} [ZW + YW]^p + \mathbb{E} Z^{2p}). \quad (3.8)$$

This lemma will be proved shortly. We first show how Proposition 3.2 follows from this lemma. By the Burkholder-Davis-Gundy inequalities, we obtain

$$\mathbb{E} \sup_n |M_n|^{2p} \leq c p^{4p} (\mathbb{E} [ZW + YW]^p + \mathbb{E} Z^{2p}). \quad (3.9)$$

Combining with (3.7) and the fact that  $Q_m = M_m - V_m$  and then using Cauchy-Schwarz completes the proof of Proposition 3.2 in the special case where the  $Q^i$  are constant from some  $n_0$  on. In the general case, let  $\bar{Q}_n^i = Q_{n \wedge n_0}^i$  for  $i = 1, 2$ , obtain (3.5) for  $\bar{Q}_n = \bar{Q}_n^1 - \bar{Q}_n^2$ , let  $n_0 \rightarrow \infty$ , and apply monotone convergence.  $\square$

**Proof of Lemma 3.3.** We now prove (3.8). Simple algebraic manipulations show that

$$V_{\infty}^2 - V_m^2 = \sum_{n=m}^{\infty} (V_{n+1} - V_n)^2 + 2 \sum_{n=m}^{\infty} V_n (V_{n+1} - V_n). \quad (3.10)$$

(Note that the sums are actually finite because  $V_m = 0$  if  $m \geq n_0$ .) Recalling  $Q = M - V$  and  $V_{\infty} = 0$ , we have

$$\begin{aligned} \sum_{n=m}^{\infty} (M_{n+1} - M_n)^2 &\leq 2 \sum_{n=m}^{\infty} (V_{n+1} - V_n)^2 + 2 \sum_{n=m}^{\infty} (Q_{n+1} - Q_n)^2 \\ &= -2V_m^2 - 4 \sum_{n=m}^{\infty} V_n (V_{n+1} - V_n) + 2 \sum_{n=m}^{\infty} (Q_{n+1} - Q_n)^2 \\ &\leq -4 \sum_{n=m}^{\infty} V_n (V_{n+1} - V_n) + 2 \sum_{n=m}^{\infty} (Q_{n+1} - Q_n)^2 \\ &=: -4S_1 + 2S_2. \end{aligned} \quad (3.11)$$

We now take the conditional expectation with respect to  $\mathcal{F}_m$ .

$$\begin{aligned}
\mathbb{E}[S_1 | \mathcal{F}_m] &= \mathbb{E}\left[\sum_{n=m}^{\infty} V_n(V_{n+1} - V_n) | \mathcal{F}_m\right] \\
&= \mathbb{E}\left[\sum_{n=m}^{\infty} \mathbb{E}[V_{n+1} - V_n | \mathcal{F}_n] V_n | \mathcal{F}_m\right] \\
&= -\mathbb{E}\left[\sum_{n=m}^{\infty} \mathbb{E}[Q_{n+1} - Q_n | \mathcal{F}_n] V_n | \mathcal{F}_m\right] \\
&= -\mathbb{E}\left[\sum_{n=m}^{\infty} V_n(Q_{n+1} - Q_n) | \mathcal{F}_m\right] \\
&= -\mathbb{E}\left[\sum_{n=m}^{\infty} V_{n+1}(Q_{n+1} - Q_n) | \mathcal{F}_m\right] \\
&\quad + \mathbb{E}\left[\sum_{n=m}^{\infty} (V_{n+1} - V_n)(Q_{n+1} - Q_n) | \mathcal{F}_m\right] \\
&=: I_1 + I_2.
\end{aligned}$$

Since  $V_{n+1} \leq 2\mathbb{E}[Z | \mathcal{F}_{n+1}]$  by (3.6), and recalling that  $Q_n = Q_n^1 - Q_n^2$ , we have

$$|I_1| \leq 2\mathbb{E}\left[\sum_{n=m}^{\infty} Z|Q_{n+1} - Q_n| | \mathcal{F}_m\right] \leq 2\mathbb{E}[ZW | \mathcal{F}_m].$$

Recalling that  $V_n = M_n - Q_n$  and setting  $S_3 = \sum_{n=m}^{\infty} (M_{n+1} - M_n)(Q_{n+1} - Q_n)$  we see that

$$I_2 = -\mathbb{E}[S_2 | \mathcal{F}_m] + \mathbb{E}[S_3 | \mathcal{F}_m].$$

Since

$$\sum_{n=m}^{\infty} (Q_{n+1} - Q_n)^2 \leq YW \tag{3.12}$$

we have

$$\mathbb{E}[S_2 | \mathcal{F}_m] \leq \mathbb{E}[YW | \mathcal{F}_m].$$

Let  $J = \mathbb{E}[\sum_{n=m}^{\infty} (M_{n+1} - M_n)^2 | \mathcal{F}_m]$ . By Cauchy-Schwarz and (3.12),

$$|\mathbb{E}[S_3 | \mathcal{F}_m]| \leq J^{1/2} \left(\mathbb{E}[YW | \mathcal{F}_m]\right)^{1/2}.$$

We therefore conclude

$$J \leq c\mathbb{E}[ZW + YW | \mathcal{F}_m] + c_1 J^{1/2} \left(\mathbb{E}[YW | \mathcal{F}_m]\right)^{1/2}. \tag{3.13}$$

Using the inequality  $A^{1/2}x^{1/2} \leq (A+x)/2$  with  $x = J$  and  $A = c_1^2 \mathbb{E}[YW \mid \mathcal{F}_m]$ , we see that

$$J \leq c \mathbb{E}[ZW + YW \mid \mathcal{F}_m] + c_1^2 \left( \mathbb{E}[YW \mid \mathcal{F}_m] \right) / 2 + J/2 \quad (3.14)$$

and therefore

$$J = \mathbb{E} \left[ \sum_{n=m}^{\infty} (M_{n+1} - M_n)^2 \mid \mathcal{F}_m \right] \leq c \mathbb{E}[ZW + YW \mid \mathcal{F}_m]. \quad (3.15)$$

We have  $|Q_{n+1} - Q_n| \leq Y$  for all  $n$  and so

$$\mathbb{E} \sup_n |Q_{n+1} - Q_n|^p \leq \mathbb{E} Y^p. \quad (3.16)$$

Using (3.7), (3.16) and the fact that  $Q_m = M_m - V_m$  and  $Y \leq (YW)^{1/2}$  we then have that

$$\begin{aligned} \mathbb{E} \sup_n |M_{n+1} - M_n|^{2p} &\leq c^p \mathbb{E} \sup_n |Q_{n+1} - Q_n|^{2p} + c^p \mathbb{E} \sup_n |V_{n+1} - V_n|^{2p} \\ &\leq c^p (\mathbb{E} Z^{2p} + \mathbb{E} (YW)^p). \end{aligned} \quad (3.17)$$

(3.8) then follows using (3.15) and Proposition 3.1 with  $A_n = \sum_{j=1}^n (M_j - M_{j-1})^2$ .  $\square$

#### 4. The $k = 2$ case.

**Proposition 4.1.** *If*

$$\tilde{U}_2(n, x) = \sum_{i=0}^{n-1} G_n(X_n - X_i - x),$$

then

$$M_n = \tilde{U}_2(n, x) + \tilde{B}_2(n, x)$$

is a martingale with  $M_0 = 0$ .

**Proof.** If

$$U_2(n, x) = \sum_{i=0}^{n-1} G(X_n - X_i - x),$$

we have  $\tilde{U}_2(n, x) = U_2(n, x) - nG(e_1\sqrt{n})$  so that

$$M_n = U_2(n, x) + B_2(n, x) - nG(x).$$

Abbreviating  $\bar{B}_n = B_2(n, x) - nG(x)$  we have

$$\bar{B}_n - \bar{B}_{n-1} = \sum_{i=0}^{n-1} 1_{(X_n = X_i + x)} - G(x).$$

So

$$\begin{aligned}
\mathbb{E}[\bar{B}_n - \bar{B}_{n-1} + G(x) \mid \mathcal{F}_{n-1}] &= \sum_{i=0}^{n-1} \mathbb{P}(X_n - X_{n-1} + X_{n-1} - X_i = x \mid \mathcal{F}_{n-1}) \\
&= \sum_{i=0}^{n-1} p(1, 0, X_{n-1} - X_i - x)
\end{aligned} \tag{4.1}$$

Abbreviating  $U_n = U_2(n, x)$  we have

$$U_n - U_{n-1} - G(x) = \sum_{i=0}^{n-1} [G(X_n - X_i - x) - G(X_{n-1} - X_i - x)].$$

Now for any  $i \leq n-1$

$$\begin{aligned}
\mathbb{E}[G(X_n - X_i - x) \mid \mathcal{F}_{n-1}] &= \mathbb{E}[G(X_n - X_{n-1} + X_{n-1} - X_i - x) \mid \mathcal{F}_{n-1}] \\
&= \sum_y G(y + X_{n-1} - X_i - x) \mathbb{P}(X_n - X_{n-1} = y \mid \mathcal{F}_{n-1}) \\
&= \sum_y G(y + X_{n-1} - X_i - x) \mathbb{P}(X_n - X_{n-1} = y) \\
&= \sum_y G(y + X_{n-1} - X_i - x) p(1, 0, y) \\
&= P_1 G(X_{n-1} - X_i - x)
\end{aligned} \tag{4.2}$$

where  $P_j$  is the transition operator associated to  $p(j, x, y)$ . Hence

$$\mathbb{E}[U_n - U_{n-1} - G(x) \mid \mathcal{F}_{n-1}] = \sum_{i=0}^{n-1} [P_1 G(X_{n-1} - X_i - x) - G(X_{n-1} - X_i - x)].$$

Comparing with (4.1) and using

$$P_1 G(z) - G(z) = -p(1, 0, z), \quad z \in \mathbb{Z}^2, \tag{4.3}$$

we see that

$$\mathbb{E}[M_n - M_{n-1} \mid \mathcal{F}_{n-1}] = \mathbb{E}[U_n - U_{n-1} + \bar{B}_n - \bar{B}_{n-1} \mid \mathcal{F}_{n-1}] = 0$$

as required. □

The key to proving Theorem 1.1 in the  $k = 2$  case is the following proposition.

**Proposition 4.2.** *We have*

$$\mathbb{E} |\tilde{\beta}_2(n, x) - \tilde{\beta}_2(n, x')|^p \leq c(p)(\log n)^p n |x - x'|^{p/3} \quad (4.4)$$

for each integer  $p > 1$  and  $x, x' \in \mathbb{Z}^2/\sqrt{n}$  with  $|x|, |x'| \leq 1$ .

Let

$$W_2(n) = |B_2(n, x)| + |B_2(n, x')|,$$

$$Y_2(n) = \max_{i \leq n} \{|B_2(i, x) - B_2(i-1, x)| + |B_2(i, x') - B_2(i-1, x')|\},$$

and

$$\tilde{Z}_2(n) = \sup_{j \leq n} |\tilde{U}_2(j, x) - \tilde{U}_2(j, x')|.$$

In the proof of Proposition 4.2 we will need the following three lemmas, whose proofs are deferred until the next section.

**Lemma 4.3.** *For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$*

$$\mathbb{E} W_2(n)^p \leq c(\log n)^p n^p. \quad (4.5)$$

**Lemma 4.4.** *For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$*

$$\mathbb{E} Y_2(n)^p \leq cn(\log n)^p. \quad (4.6)$$

**Lemma 4.5.** *For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$*

$$\mathbb{E} \tilde{Z}_2(n)^p \leq cn^p \left| \frac{x - x'}{\sqrt{n}} \right|^{2p/3}, \quad (4.7)$$

**Proof of Proposition 4.2.** Converting from  $\tilde{\beta}$ 's to  $\tilde{B}$ 's, estimate (4.4) for  $k = 2$  is equivalent to

$$\mathbb{E} |\tilde{B}_2(n, x) - \tilde{B}_2(n, x')|^p \leq c(p)(\log n)^p n^{p+1} \left( \frac{|x - x'|}{\sqrt{n}} \right)^{p/3} \quad (4.8)$$

for  $x, x' \in \mathbb{Z}^2$  with  $|x|, |x'| \leq \sqrt{n}$ . We want to apply Proposition 3.2. We fix an  $n$ . We use the notation  $f^+(x) = \max(f(x), 0)$ ,  $f^-(x) = \max(-f(x), 0)$  so that  $f(x) = f^+(x) - f^-(x)$ . Take for  $i \leq n$

$$Q_i^1 = B_2(i, x) + (G^+(x') + G^-(x))i, \quad Q_i^2 = B_2(i, x') + (G^+(x) + G^-(x'))i, \quad (4.9)$$

so that  $Q^1$  and  $Q^2$  are increasing and  $Q_i = Q_i^1 - Q_i^2 = \tilde{B}_2(i, x) - \tilde{B}_2(i, x')$ . For  $i \geq n$  and  $j = 1, 2$ , set  $Q_i^j = Q_n^j$ . We set  $H_i = \tilde{U}_2(i, x) - \tilde{U}_2(i, x')$ . By Proposition 4.1,  $Q_i + H_i$  is a martingale.

From Proposition 2.1, Lemmas 4.3 and 4.4, and the fact that  $|x|, |x'| \leq \sqrt{n}$ , we see that

$$\mathbb{E}[(Q_n^1 + Q_n^2)^p] \leq c(\log n)^p n^p \quad (4.10)$$

and

$$\mathbb{E}[(\max_{i \leq n} \{[Q_i^1 - Q_{i-1}^1] + [Q_i^2 - Q_{i-1}^2]\})^p] \leq cn(\log n)^p. \quad (4.11)$$

Combining (4.10), (4.11), Lemma 4.5, and the fact that  $\frac{1}{\sqrt{n}} \leq \frac{|x-x'|}{\sqrt{n}} \leq 2$  unless  $x = x'$  with Proposition 3.2, we obtain

$$\mathbb{E} \sup_{j \leq n} |\tilde{B}_2(j, x) - \tilde{B}_2(j, x')|^p \leq c(p)(\log n)^p n^{p+1} \left( \frac{|x-x'|}{\sqrt{n}} \right)^{p/3} \quad (4.12)$$

for  $x, x' \in \mathbb{Z}^2$  with  $|x|, |x'| \leq \sqrt{n}$ , which implies (4.8). This is the bound we need.  $\square$

**Proof of Theorem 1.1, the  $k = 2$  case.** Let  $f : \mathbb{R}^2 \rightarrow [0, \infty)$  be a nonnegative  $C^\infty$  function with support in  $\{y : \frac{1}{2} \leq |y| \leq 1\}$ , and with integral 1. Let  $f_\tau(x) = \tau^{-2} f(x/\tau)$ . The gradient of  $f_\tau$  is bounded by a constant times  $\tau^{-3}$ . Set  $\tau_n = n^{-\zeta/4}$ . Then recalling (1.7),

$$\int_0^1 \int_0^t |f_{\tau_n}(X_t^n - X_s^n) - f_{\tau_n}(W_t^n - W_s^n)| ds dt \leq c\tau_n^{-3} n^{-\zeta} \leq cn^{-\zeta/4}. \quad (4.13)$$

We also have by Lemma 2.3 that for some  $\bar{\delta} > 0$

$$\left| \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f_{\tau_n}(x) G_n(x\sqrt{n}) \frac{1}{n} - \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f_{\tau_n}(x) \frac{1}{\pi} \log(1/|x|) \frac{1}{n} \right| \leq cn^{-\bar{\delta}} \quad (4.14)$$

and it is easy to see from the support properties of  $f_{\tau_n}(x)$  that

$$\left| \int f_{\tau_n}(x) \frac{1}{\pi} \log(1/|x|) dx - \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f_{\tau_n}(x) \frac{1}{\pi} \log(1/|x|) \frac{1}{n} \right| \leq cn^{-\bar{\delta}}. \quad (4.15)$$

On the other hand, recalling the notation  $X_t^n = X_{[nt]}/\sqrt{n}$

$$\begin{aligned} \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f_{\tau_n}(x) B_2(n, \sqrt{n}x) \frac{1}{n^2} &= \frac{1}{n^2} \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f_{\tau_n}(x) \sum_{0 \leq i_1 < i_2 \leq n} 1_{(X_{i_2} = X_{i_1} + \sqrt{n}x)} \\ &= \frac{1}{n^2} \sum_{0 \leq i_1 < i_2 \leq n} f_{\tau_n} \left( \frac{(X_{i_2} - X_{i_1})}{\sqrt{n}} \right) \\ &= \frac{1}{n^2} \int_0^n \int_0^t f_{\tau_n} \left( \frac{(X_{[t]} - X_{[s]})}{\sqrt{n}} \right) ds dt \\ &= \int_0^1 \int_0^t f_{\tau_n}(X_t^n - X_s^n) ds dt \end{aligned} \quad (4.16)$$



so that

$$\sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \tilde{\beta}_2(n, x) \frac{1}{n} = \int_0^1 \int_0^t f_{\tau_n}(X_t^n - X_s^n) ds dt - \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) G_n(x\sqrt{n}) \frac{1}{n}.$$

By [3],

$$\int f_{\tau_n}(x) \tilde{\gamma}_2(1, x, n) dx = \int_0^1 \int_0^1 f_{\tau_n}(W_t^n - W_s^n) ds dt - \int f_{\tau_n}(x) \frac{1}{\pi} \log(1/|x|) dx. \quad (4.17)$$

(This conforms with the definition given in Section 1 above; the definition in [3] is very slightly different and would yield  $\int_0^1 \int_0^1 f_{\tau_n}(W_t^n - W_s^n) ds dt$  instead.) Combining the above,

$$\left| \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \tilde{\beta}_2(n, x) \frac{1}{n} - \int f_{\tau_n}(x) \tilde{\gamma}_2(1, x, n) dx \right| = O(n^{-\zeta/4}) + O(n^{-\bar{\delta}}). \quad (4.18)$$

Recall  $\int f_{\tau_n}(x) dx = 1$ . Without loss of generality we may assume  $\zeta < 1/2$  is small enough so that  $\psi_n =: \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \frac{1}{n} = 1 + O(n^{-\bar{\delta}})$  for some  $\bar{\delta} > 1/4$ . (If  $\zeta$  were too large, then  $\tau_n$  would tend to 0 too quickly, and then the above estimate for  $\psi_n$  might not be valid. In general one has  $\psi_n = 1 + O(n^{-1/2} \tau_n^{-3})$ .) Jensen's inequality and estimates (4.4), (4.5) imply that

$$\begin{aligned} \mathbb{E} \left| \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \tilde{\beta}_2(n, x) \frac{1}{n} - \psi_n \tilde{\beta}_2(n, 0) \right|^p & \quad (4.19) \\ & \leq c(p) \mathbb{E} \left| \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} \frac{1}{n} f_{\tau_n}(x) [\tilde{\beta}_2(n, x) - \tilde{\beta}_2(n, 0)] \right|^p + c(p) (\psi_n - 1)^p \mathbb{E} \tilde{\beta}_2(n, 0)^p \\ & \leq c(p) \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \mathbb{E} |\tilde{\beta}_2(n, x) - \tilde{\beta}_2(n, 0)|^p \frac{1}{n} + c(p) n^{-p\bar{\delta}} (\log n)^p \\ & \leq c(p) (\log n)^p n (\tau_n)^{p/3}. \end{aligned}$$

If we take  $p$  big enough, then using Chebyshev's inequality

$$\mathbb{P} \left( \left| \sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \tilde{\beta}_2(n, x) \frac{1}{n} - \psi_n \tilde{\beta}_2(n, 0) \right| \geq n^{-\zeta/24} \right) \leq c \frac{(\log n)^p n (\tau_n)^{p/3}}{n^{-\zeta p/24}} \leq \frac{c}{n^2}.$$

By Borel-Cantelli, we conclude that

$$\sum_{x \in \mathbb{Z}^2 / \sqrt{n}} f_{\tau_n}(x) \tilde{\beta}_2(n, x) \frac{1}{n} - \psi_n \tilde{\beta}_2(n, 0) = O(n^{-\zeta/24}), \quad a.s. \quad (4.20)$$

Using (4.10),

$$\mathbb{E}[|\tilde{B}_2(n, 0)^p|] \leq c(\log n)^p n^p,$$

so

$$\mathbb{P}(|\tilde{\beta}_2(n, 0)| > n^{\bar{\delta}/2}) \leq \frac{c(\log n)^p}{n^{\bar{\delta}p/2}},$$

and if we take  $p$  large enough, Borel-Cantelli tells us that

$$\tilde{\beta}_2(n, 0) = O(n^{\bar{\delta}/2}), \quad a.s.$$

So then

$$\tilde{\beta}_2(n, 0) - \psi_n \tilde{\beta}_2(n, 0) = O(n^{-\bar{\delta}/2}), \quad a.s. \quad (4.21)$$

A very similar argument to the above also shows that we have

$$\int f_{\tau_n}(x) \tilde{\gamma}_2(1, x, n) dx - \tilde{\gamma}_2(1, 0, n) = O(n^{-\zeta/24}), \quad a.s. \quad (4.22)$$

the analogue to estimate (4.4) is in [3]. Combining (4.18), (4.20), (4.21) and (4.22) we conclude that

$$\tilde{\beta}_2(n, 0) - \tilde{\gamma}_2(1, 0, n) = O(n^{-\zeta/24}) + O(n^{-\bar{\delta}/2}), \quad a.s.$$

□

**Remark 4.6.** To see the importance of renormalization, note that if we also had the estimate (4.8) for  $B_2(n, x) - B_2(n, x')$ , this would imply that uniformly in  $n$

$$|G(x) - G(x')| \leq c(p)(\log n) n^{1/p} \left( \frac{|x - x'|}{\sqrt{n}} \right)^{1/3}$$

which is impossible if  $p > 6$  and  $n$  is sufficiently large.

## 5. Proofs of Lemmas 4.3-4.5.

**Proof of Lemma 4.3.** We have

$$\begin{aligned} & \mathbb{E}\{(B_2(n, x))^p\} \\ & \leq \mathbb{E}\left\{\left(\sum_{i,j=0}^n 1_{(X_j - X_i = x)}\right)^p\right\} \\ & = \sum_{s \in \mathcal{S}} \sum_{0 \leq i_1 \leq \dots \leq i_{2p} \leq n} \sum_{z_1, \dots, z_p \in \mathbb{Z}^2} \prod_{j=1}^{2p} p(i_j - i_{j-1}, z_{s(j-1)} + x_{c(j-1)}, z_{s(j)} + x_{c(j)}) \end{aligned} \quad (5.1)$$

where  $s$  runs over the set of maps  $\mathcal{S}$  from  $\{1, \dots, 2p\}$  to  $\{1, \dots, p\}$  such that  $s^{-1}(j) = 2$  for each  $1 \leq j \leq p$ ,  $c(j) = \sum_{i=1}^j 1_{\{s(i)=s(j)\}}$  and  $x_1 = 0, x_2 = x$ . Here we use the conventions  $i_0 = 0, z_0 = 0, c(0) = 0$ . Setting

$$g_n(x) = \sum_{i=0}^n p(i, 0, x) \leq c \log n \quad (5.2)$$

for  $x \leq \sqrt{n}$  by Proposition 2.1, and using the obvious fact that

$$\sum_{x \in \mathbb{Z}^2} g_n(x) = \sum_{x \in \mathbb{Z}^2} \sum_{i=0}^n p(i, 0, x) = n + 1$$

we can bound (5.1) by

$$\begin{aligned} & \sum_{s \in \mathcal{S}} \sum_{z_1, \dots, z_p \in \mathbb{Z}^2} \prod_{j=1}^{2p} g_n(z_{s(j)} + x_{c(j)} - z_{s(j-1)} - x_{c(j-1)}) \\ & \leq c(\log n)^p \sum_{s \in \mathcal{S}} \sum_{z_1, \dots, z_p \in \mathbb{Z}^2} \prod_{j: c(j)=1} g_n(z_{s(j)} + x_{c(j)} - z_{s(j-1)} - x_{c(j-1)}) \\ & \leq c(n \log n)^p. \end{aligned} \quad (5.3)$$

□

**Proof of Lemma 4.4.** Let

$$C(m, x) = [B_2(m, x) - B_2(m-1, x)] = \sum_{i=0}^{m-1} 1_{(X_m = X_i + x)}.$$

We show that for any  $1 \leq m \leq n$

$$\mathbb{E} C(m, x)^p \leq c(\log n)^p; \quad (5.4)$$

using

$$Y_2(n)^p \leq \sum_{m=1}^n c(p) C(m, x)^p,$$

we are then done.

But

$$\begin{aligned}
& \mathbb{E} \left\{ \left( \sum_{i=0}^{m-1} 1_{(X_m = X_i + x)} \right)^p \right\} \\
& \leq p! \sum_{0 \leq i_1 \leq \dots \leq i_p \leq m-1} \sum_{z_1, \dots, z_p, y \in \mathbb{Z}^2} \prod_{j=1}^p p(i_j - i_{j-1}, z_{j-1}, z_j) 1_{(y = z_j + x)} p(m - i_p, z_p, y) \\
& = p! \sum_{0 \leq i_1 \leq \dots \leq i_p \leq m-1} \sum_{y \in \mathbb{Z}^2} p(i_1, 0, y - x) \prod_{j=2}^p p(i_j - i_{j-1}, 0, 0) p(m - i_p, 0, x) \\
& = p! \sum_{0 \leq i_1 \leq \dots \leq i_p \leq m-1} \prod_{j=2}^p p(i_j - i_{j-1}, 0, 0) p(m - i_p, 0, x) \\
& \leq c(\log m)^p
\end{aligned}$$

which gives (5.4). □

**Proof of Lemma 4.5.** We begin by estimating

$$\mathbb{E} (1 + |X_i|^2)^{-b/2}$$

with  $0 < b < 2$ .

First

$$\mathbb{E} [(1 + |X_i|^2)^{-b/2}; |X_i| = 0] = 1 \cdot \mathbb{P}(X_i = 0) \leq c/i \leq ci^{-b/2}.$$

Next, using (2.1)

$$\begin{aligned}
\mathbb{E} [(1 + |X_i|^2)^{-b/2}; 0 < |X_i| < \sqrt{i}] & \leq \sum_{\{x \in \mathbb{Z}^2, 0 < |x| \leq \sqrt{i}\}} |x|^{-b} p(i, 0, x) \\
& \leq \frac{c}{i} \sum_{\{x \in \mathbb{Z}^2, 0 < |x| \leq \sqrt{i}\}} |x|^{-b} \\
& = \frac{c}{i} i^{1-b/2} = ci^{-b/2}.
\end{aligned}$$

Finally,

$$\mathbb{E} [(1 + |X_i|^2)^{-b/2}; \sqrt{i} \leq |X_i|] \leq (1 + i)^{-b/2} \mathbb{P}(\sqrt{i} \leq |X_i|) \leq ci^{-b/2}. \quad (5.5)$$

We conclude that for any  $0 < b < 2$

$$\mathbb{E} [(1 + |X_i|^2)^{-b/2}] \leq ci^{-b/2}. \quad (5.6)$$

Using the estimate

$$|G(X_i + x) - G(X_i + y)| \leq \frac{c|x - y|^{2/3}}{(1 + |X_i + x|^2)^{1/3}} + \frac{c|x - y|^{2/3}}{(1 + |X_i + y|^2)^{1/3}}$$

of Proposition 2.2, the fact that symmetry tells us that  $\mathbb{E}[(1 + |X_i + x|^2)^{-1/3}]$  is largest when  $x = 0$ , and the estimate (5.6) above, we obtain

$$\mathbb{E} \sum_{i=1}^n |G(X_i + x) - G(X_i + y)| \leq c|x - y|^{2/3} \sum_{i=1}^n i^{-1/3} \leq cn(|x - y|/\sqrt{n})^{2/3}.$$

So by independence, using  $\bar{X}_i, \bar{\mathbb{E}}$  to denote an independent copy of  $X_i$  and its expectation operator,

$$\begin{aligned} & \mathbb{E} \left[ \sum_{i=m+1}^n |G(X_i + x) - G(X_i + y)| \mid \mathcal{F}_m \right] \\ & \leq \bar{\mathbb{E}} \sum_{i=1}^n |G(\bar{X}_i + X_m + x) - G(\bar{X}_i + X_m + y)| \leq cn \left( \frac{|x - y|}{\sqrt{n}} \right)^{2/3}. \end{aligned} \quad (5.7)$$

If  $|x| \leq \sqrt{n}$ , by Proposition 2.1 and Doob's inequality

$$\begin{aligned} \mathbb{E} \sup_{i \leq n} |G(X_i + x)|^p & \leq c \mathbb{E} \sup_{i \leq n} (1 + \log^+ |X_i + x|)^p \\ & \leq c(\log n)^p \mathbb{P}(\sup_{i \leq n} |X_i + x| \leq n) + c \sum_{m=\lceil \log n \rceil}^{\infty} m^p \mathbb{P}(e^m \leq \sup_{i \leq n} |X_i + x| \leq e^{m+1}) \\ & \leq c(\log n)^p + c \sum_{m=\lceil \log n \rceil}^{\infty} m^p \frac{\mathbb{E} |X_n + x|^2}{e^{2m}} \\ & \leq c(\log n)^p. \end{aligned} \quad (5.8)$$

If  $x \neq y$ , then  $|x - y| \geq 1$  and (5.8) then implies that

$$\mathbb{E} \sup_{i \leq n} |G(X_i + x) - G(X_i + y)|^p \leq c(\log n)^p \leq c \left( n \left( \frac{|x - y|}{\sqrt{n}} \right)^{2/3} \right)^p.$$

Using Proposition 3.1 with  $A_j = \sum_{i=1}^{j \wedge n} |G(X_i + x) - G(X_i + y)|$ , if  $|x|, |y| \leq \sqrt{n}$ , then

$$\left\| \sum_{i=1}^n |G(X_i + x) - G(X_i + y)| \right\|_p \leq cn(|x - y|/\sqrt{n})^{2/3}. \quad (5.9)$$

Replacing  $x$  and  $y$  by  $-x$  and  $-x'$ , resp., and using the fact that  $\sum_{i=1}^n G(X_i - x)$  is equal in law to  $\sum_{i=0}^{n-1} G(X_n - X_i - x)$  yields the  $L^p$  estimate that we want.  $\square$

**6. The martingale connection:**  $k > 2$ .

Let  $\tilde{B}_{1,m}(j, x) = j$  and for  $k \geq 2$  define

$$\tilde{B}_{k,m}(j, x) = \sum_{A \subset \{2, \dots, k\}} (-1)^{|A|} \left( \prod_{i \in A} G_m(x_i) \right) B_{k-|A|}(j, x_{A^c}). \quad (6.1)$$

Note that  $\tilde{B}_{k,n}(n, x) = \tilde{B}_k(n, x)$ .

If  $x = (x_2, \dots, x_{k-1}, x_k)$ , let  $x_{k^c} = (x_2, \dots, x_{k-1})$ .

**Proposition 6.1.** *Let  $k > 2$ . If*

$$\tilde{U}_{k,m}(n, x) = \sum_{i=1}^n G_m(X_n - X_i - x_k) [\tilde{B}_{k-1,m}(i, x_{k^c}) - \tilde{B}_{k-1,m}(i-1, x_{k^c})],$$

then for each  $m$

$$M_{n,m} = \tilde{U}_{k,m}(n, x) + \tilde{B}_{k,m}(n, x), \quad n = 0, 1, 2, \dots$$

is a martingale with  $M_{0,m} = 0$ .

**Proof.** We will show that for each  $k$

$$N_{k,m}(n) = U_{k,m}(n, x) + B_{k,m}(n, x) - G_m(x_k) B_{k-1,m}(n, x_{k^c}), \quad n = 0, 1, 2, \dots$$

is a martingale where

$$\begin{aligned} U_{k,m}(n, x) &= \sum_{i=1}^n [G(X_n - X_i - x_k) - G(e_1 \sqrt{m})] [B_{k-1}(i, x_{k^c}) - B_{k-1}(i-1, x_{k^c})] \\ &= \sum_{i=1}^n G(X_n - X_i - x_k) [B_{k-1}(i, x_{k^c}) - B_{k-1}(i-1, x_{k^c})] \\ &\quad - G(e_1 \sqrt{m}) B_{k-1}(n, x_{k^c}). \end{aligned}$$

This will prove the proposition since, with the notation  $D_k = D \cup \{k\}$ ,

$$\begin{aligned} &\tilde{U}_{k,m}(n, x) + \tilde{B}_{k,m}(n, x) \\ &= \sum_{i=1}^n G_m(X_n - X_i - x_k) \\ &\quad \sum_{D \subset \{2, \dots, k-1\}} (-1)^{|D|} \left( \prod_{l \in D} G_m(x_l) \right) [B_{k-|D|,m}(i, x_{D_k^c}) - B_{k-|D|,m}(i-1, x_{D_k^c})] \\ &\quad + \sum_{A \subset \{2, \dots, k\}} (-1)^{|A|} \left( \prod_{i \in A} G_m(x_i) \right) B_{k-|A|}(n, x_{A^c}) \\ &= \sum_{D \subset \{2, \dots, k-1\}} (-1)^{|D|} \left( \prod_{l \in D} G_m(x_l) \right) \\ &\quad [U_{k-|D^c|,m}(n, x_{D^c}) + B_{k-|D^c|,m}(n, x_{D^c}) - G_m(x_k) B_{k-|D^c|-1,m}(n, x_{D_k^c})]. \end{aligned}$$

If we set

$$\bar{U}_k(n, x) = \sum_{i=1}^n G(X_n - X_i - x_k)[B_{k-1}(i, x_{k^c}) - B_{k-1}(i-1, x_{k^c})]$$

we have that

$$N_{k,m}(n) = \bar{U}_k(n, x) + B_k(n, x) - G(x_k)B_{k-1}(n, x_{k^c}).$$

Abbreviating  $\bar{U}_n = \bar{U}_k(n, x)$  and  $\bar{B}_n = B_k(n, x) - G(x_k)B_{k-1}(n, x_{k^c})$ , we have that

$$N_{k,m}(n) = \bar{U}_n + \bar{B}_n.$$

Setting

$$H_i = B_{k-1}(i, x_{k^c}) - B_{k-1}(i-1, x_{k^c}) = \sum_{0 \leq i_1 < i_2 < \dots < i_{k-1} = i} \prod_{j=2}^{k-1} 1_{(X_{i_j} = X_{i_{j-1}} + x_j)}$$

we have

$$\bar{B}_n - \bar{B}_{n-1} = \sum_{i=1}^{n-1} 1_{(X_n = X_i + x_k)} H_i - G(x_k)H_n.$$

So using (4.1)

$$\begin{aligned} & \mathbb{E}[\bar{B}_n - \bar{B}_{n-1} + G(x_k)H_n \mid \mathcal{F}_{n-1}] \\ &= \sum_{i=1}^{n-1} p(1, 0, X_{n-1} - X_i - x_k) H_i. \end{aligned} \tag{6.2}$$

From the definition of  $\bar{U}_n$  we have

$$\begin{aligned} \bar{U}_n - \bar{U}_{n-1} &= G(x_k)H_n \\ &+ \sum_{i=1}^{n-1} [G(X_n - X_i - x_k) - G(X_{n-1} - X_i - x_k)] H_i. \end{aligned}$$

Recalling (4.2)

$$\begin{aligned} & \mathbb{E}[\bar{U}_n - \bar{U}_{n-1} - G_n(x_k)H_n \mid \mathcal{F}_{n-1}] \\ &= \sum_{i=1}^{n-1} [P_1 G(X_{n-1} - X_i - x_k) - G(X_{n-1} - X_i - x_k)] H_i. \end{aligned}$$

Comparing with (6.2) and using (4.3)

$$P_1 G(x) - G(x) = -p(1, 0, x),$$

we see that

$$\mathbb{E}[\bar{U}_n - \bar{U}_{n-1} + \bar{B}_n - \bar{B}_{n-1} \mid \mathcal{F}_{n-1}] = 0$$

as required.  $\square$

**Remark.** The statement of Proposition 6.1 is not an exact analogue of that of Proposition 4.1. Consider the summands in the definition of  $\tilde{U}_{k,m}(n, x)$ :

$$G_m(X_n - X_i - x_k)[\tilde{B}_{k-1,m}(i, x_{k^c}) - \tilde{B}_{k-1,m}(i-1, x_{k^c})]. \quad (6.3)$$

When  $k = 2$  and  $i = n$ , this is nonrandom, whereas this is not the case when  $k > 2$  and  $i = n$ . On the other hand, recalling that  $B_{k-1}(i, x) = 0$  if  $i > k-1$ , it is natural to define  $B_{k-1}(-1, x)$  to be 0. It is also natural to define  $\tilde{B}_{1,m}(i, x) = i$  for  $i \geq 0$ . Then (6.3) will be 0 if  $i = 0$  for all  $k \geq 2$ , but the  $i = 0$  term in the statement of Proposition 4.1 is not zero.

## 7. The case of general $k$ .

The key to proving Theorem 1.1 for the case of general  $k$  is the following proposition.

**Proposition 7.1.** *For any  $k \geq 2$  we have*

$$\mathbb{E}|\tilde{\beta}_k(n, x) - \tilde{\beta}_k(n, x')|^p \leq c(p)(\log n)^{p(k-1)}n^k|x - x'|^{8^{-k}p} \quad (7.1)$$

for each integer  $p > 1$  and  $x, x' \in (\mathbb{Z}^2)^{k-1}/\sqrt{n}$  with  $|x|, |x'| \leq 1$ .

Let

$$W_k(n) = |B_k(n, x)| + |B_k(n, x')|,$$

$$\tilde{W}_k(n) = |\tilde{B}_k(n, x)| + |\tilde{B}_k(n, x')|,$$

$$Y_k(n) = \max_{i \leq n} \{|B_k(i, x) - B_k(i-1, x)| + |B_k(i, x') - B_k(i-1, x')|\},$$

$$\tilde{Y}_k(n) = \max_{i \leq n} \{|\tilde{B}_{k,n}(i, x) - \tilde{B}_{k,n}(i-1, x)| + |\tilde{B}_{k,n}(i, x') - \tilde{B}_{k,n}(i-1, x')|\},$$

and

$$\tilde{Z}_k(n) = \sup_{j \leq n} |\tilde{U}_{k,n}(j, x) - \tilde{U}_{k,n}(j, x')|.$$

In the proof of Proposition 7.1 we will need the following three lemmas, whose proofs are deferred until the next two sections.



**Lemma 7.2.** For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$

$$\mathbb{E} W_k(n)^p \leq c(\log n)^{p(k-1)} n^p \quad (7.2)$$

and

$$\mathbb{E} \widetilde{W}_k(n)^p \leq c(\log n)^{p(k-1)} n^p. \quad (7.3)$$

**Lemma 7.3.** For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$

$$\mathbb{E} Y_k(n)^p \leq cn(\log n)^{p(k-1)} \quad (7.4)$$

and

$$\mathbb{E} \widetilde{Y}_k(n)^p \leq cn(\log n)^{p(k-1)}. \quad (7.5)$$

**Lemma 7.4.** For any  $x, x'$  with  $|x|, |x'| \leq \sqrt{n}$

$$\mathbb{E} \widetilde{Z}_k(n)^p \leq c(\log n)^{p(k-1)} n^{p+k} \left( \frac{|x-x'|}{\sqrt{n}} \right)^{8^{-k} 2p}, \quad (7.6)$$

**Proof of Proposition 7.1.** Converting from  $\widetilde{\beta}$ 's to  $\widetilde{B}$ 's, estimate (7.1) is equivalent to

$$\mathbb{E} |\widetilde{B}_{k,n}(n, x) - \widetilde{B}_{k,n}(n, x')|^p \leq c(p)(\log n)^{p(k-1)} n^{p+k} \left( \frac{|x-x'|}{\sqrt{n}} \right)^{8^{-k} p} \quad (7.7)$$

for  $x, x' \in (\mathbb{Z}^2)^{k-1}$  with  $|x|, |x'| \leq \sqrt{n}$ . We want to apply Proposition 3.2. We fix an  $n$ . Let

$$\mathcal{A}_+(x) = \left\{ A \subset \{2, \dots, k\} \mid (-1)^{|A|} \prod_{i \in A} G_n(x_i) > 0 \right\}$$

$$\widetilde{B}_{k,n,+}(j, x) = \sum_{A \in \mathcal{A}_+(x)} (-1)^{|A|} \left( \prod_{i \in A} G_n(x_i) \right) B_{k-|A|}(j, x_{A^c}). \quad (7.8)$$

$$\widetilde{B}_{k,n,-}(j, x) = \sum_{A \in \mathcal{A}_+^c(x)} (-1)^{|A|} \left( \prod_{i \in A} G_n(x_i) \right) B_{k-|A|}(j, x_{A^c}). \quad (7.9)$$

For  $i \leq n$  set

$$Q_i^1 = \widetilde{B}_{k,n,+}(i, x) - \widetilde{B}_{k,n,-}(i, x'), \quad Q_i^2 = \widetilde{B}_{k,n,+}(i, x') - \widetilde{B}_{k,n,-}(i, x),$$

so that  $Q^1$  and  $Q^2$  are increasing and  $Q_i = Q_i^1 - Q_i^2 = \widetilde{B}_{k,n}(i, x) - \widetilde{B}_{k,n}(i, x')$ . For  $i \geq n$  set  $Q_i^j = Q_n^j$ ,  $j = 1, 2$ . We set  $H_i = \widetilde{U}_{k,n}(i, x) - \widetilde{U}_{k,n}(i, x')$ . By Proposition 6.1,  $Q_i + H_i$  is a martingale. Using Lemmas 7.2-7.4 and Proposition 2.1 to bound the right hand side of (3.5) in Proposition 3.2 and using the fact that  $\frac{1}{\sqrt{n}} \leq \frac{|x-x'|}{\sqrt{n}} \leq 2$  unless  $x = x'$  we obtain

$$\mathbb{E} \sup_{j \leq n} |\widetilde{B}_{k,n}(j, x) - \widetilde{B}_{k,n}(j, x')|^p \leq c(p)(\log n)^{p(k-1)} n^{p+k} \left( \frac{|x-x'|}{\sqrt{n}} \right)^{8^{-k} p} \quad (7.10)$$

for  $x, x' \in (\mathbb{Z}^2)^{k-1}$  with  $|x|, |x'| \leq \sqrt{n}$ , which implies (7.7). This is the bound we need.  $\square$

**Proof of Theorem 1.1, the general case.** The proof is quite similar to the  $k = 2$  case. Let  $f : \mathbb{R}^2 \rightarrow [0, \infty)$  be a nonnegative  $C^\infty$  function with support in  $\{y : \frac{1}{2} \leq |y| \leq 1\}$ , and with integral 1. Let  $f_\tau(x) = \tau^{-2}f(x/\tau)$ . Set  $\tau_n = n^{-\zeta/4k}$ . Set  $g_n(f) = \sum_{x \in \mathbb{Z}^2/\sqrt{n}} f(x)G_n(x\sqrt{n})\frac{1}{n}$  and  $l(f) = \frac{1}{\pi} \int f(x) \log(1/|x|)dx$ . As in (4.14), (4.15)

$$|g_n(f_{\tau_n}) - \frac{1}{\pi}l(f_{\tau_n})| \leq c(\tau_n\sqrt{n})^{-\rho}. \quad (7.11)$$

Using (1.1) and setting  $F_{\tau_n}(x_2, \dots, x_k) = \prod_{i=2}^k f_{\tau_n}(x_i)$  we have

$$\begin{aligned} \int F_{\tau_n}(x)\tilde{\gamma}_k(1, x) dx &= \sum_{j=0}^{k-1} \binom{k-1}{j} (-1)^j (l(f_{\tau_n}))^j \\ &\times \int_{0 \leq t_1 \leq \dots \leq t_{k-j} < 1} \prod_{i=2}^{k-j} f_{\tau_n}(W_{t_i} - W_{t_{i-1}}) dt_1 \cdots dt_{k-j}. \end{aligned} \quad (7.12)$$

On the other hand, as in (4.16), it is easily checked that we have

$$\begin{aligned} \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x)\tilde{\beta}_k(n, x) \frac{1}{n^{k-1}} &= \sum_{j=0}^{k-1} \binom{k-1}{j} (-1)^j (g_n(f_{\tau_n}))^j \\ &\times \int_{0 \leq t_1 \leq \dots \leq t_{k-j} < 1} \prod_{i=2}^{k-j} f_{\tau_n}(X_{t_i}^n - X_{t_{i-1}}^n) dt_1 \cdots dt_{k-j}. \end{aligned} \quad (7.13)$$

Since the gradient of  $f_\tau$  is bounded by a constant times  $\tau^{-3}$

$$\begin{aligned} \int_{0 \leq t_1 \leq \dots \leq t_{k-j} < 1} \left| \prod_{i=2}^{k-j} f_{\tau_n}(W_{t_i}^n - W_{t_{i-1}}^n) - \prod_{i=2}^{k-j} f_{\tau_n}(X_{t_i}^n - X_{t_{i-1}}^n) \right| dt_1 \cdots dt_{k-j} \\ \leq c\tau_n^{-3-2(k-j-2)}n^{-\zeta} \leq cn^{-\zeta/4}. \end{aligned} \quad (7.14)$$

Combining (7.11), (7.14), and the fact that both  $|g_n(f_{\tau_n})|$  and  $|l(f_{\tau_n})|$  are bounded by  $c \log n$  we see that

$$\begin{aligned} \left| \int F_{\tau_n}(x)\tilde{\gamma}_k(t, x, n) dx - \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x)\tilde{\beta}_k(t, x) \frac{1}{n^{k-1}} \right| \\ \leq c\tau_n^{-2(k-2)}(\tau_n\sqrt{n})^{-\rho} + c(\log n)^{k-1}n^{-\zeta/4} \leq cn^{-\zeta/8} \end{aligned} \quad (7.15)$$

if we take  $\zeta > 0$  sufficiently small.

Since  $\int F_{\tau_n}(x)dx = 1$ , we have  $\psi_{k,n} =: \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \frac{1}{n^{k-1}} = 1 + O(n^{-\bar{\delta}})$ , provided we assume, as we may without loss of generality, that  $\zeta$  is sufficiently small. Jensen's inequality and estimates (7.1), (7.2) imply that

$$\begin{aligned} & \mathbb{E} \left| \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \tilde{\beta}_k(n, x) \frac{1}{n^{k-1}} - \psi_{k,n} \tilde{\beta}_k(n, 0) \right|^p \\ & \leq c(p) \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \mathbb{E} |\tilde{\beta}_k(n, x) - \tilde{\beta}_k(n, 0)|^p \frac{1}{n^{k-1}} + c(p) (\psi_{k,n} - 1)^p \\ & \leq c(p) (\log n)^{p(k-1)} n^2 (\tau_n)^{8-k} p. \end{aligned} \quad (7.16)$$

If we take  $p$  big enough, then

$$\begin{aligned} & \mathbb{P} \left( \left| \sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \tilde{\beta}_k(n, x) \frac{1}{n^{k-1}} - \psi_{k,n} \tilde{\beta}_k(n, 0) \right| \geq n^{-8^{-k}\zeta/8k} \right) \\ & \leq c \frac{(\log n)^{p(k-1)} n^2 (\tau_n)^{8-k} p}{n^{-8^{-k}\zeta/8k}} \leq \frac{c}{n^2}. \end{aligned}$$

By Borel-Cantelli, we conclude that

$$\sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \tilde{\beta}_k(n, x) \frac{1}{n^{k-1}} - \psi_{k,n} \tilde{\beta}_k(n, 0) = O(n^{-8^{-k}\zeta/8k}), \quad a.s. \quad (7.17)$$

We use (7.3) and the same argument as in the  $k = 2$  case to show

$$\tilde{\beta}_k(n, 0) - \psi_{k,n} \tilde{\beta}_k(n, 0) = O(n^{-\bar{\delta}/2}), \quad a.s.$$

for some  $\bar{\delta} > 0$ . This with (7.17) yields

$$\sum_{x \in \mathbb{Z}^{2(k-1)}/\sqrt{n}} F_{\tau_n}(x) \tilde{\beta}_k(n, x) \frac{1}{n^{k-1}} - \tilde{\beta}_k(n, 0) = O(n^{-8^{-k}\zeta/8k}) + O(n^{-\bar{\delta}/2}), \quad a.s. \quad (7.18)$$

A similar argument shows that we have (7.18) holding with the  $\tilde{\beta}_k(n, x)$  replaced by  $\tilde{\gamma}_k(1, x, n)$ ; the analogue to estimate (7.1) is in [3]. Combining, we conclude that

$$\tilde{\beta}_k(n, 0) - \tilde{\gamma}_k(1, 0, n) = O(n^{-8^{-k}\zeta/8k}) + O(n^{-\bar{\delta}/2}), \quad a.s.$$

□

## 8. Proofs of Lemmas 7.2-7.3.

These are again similar to the  $k = 2$  case.

**Proof of Lemma 7.2.** Using Proposition 2.1 it suffices to prove (7.2) for all  $k$ .

We have

$$\begin{aligned} & \mathbb{E} \{(B_k(n, x))^m\} \\ & \leq \mathbb{E} \left\{ \left( \sum_{i_1, \dots, i_k=0}^n \prod_{j=2}^k 1_{(X_{i_j} = X_{i_{j-1}} + x_j)} \right)^m \right\} \\ & = \sum_{s \in S(k, m)} \sum_{0 \leq i_1 \leq \dots \leq i_{km} \leq n} \sum_{z_1, \dots, z_m \in \mathbb{Z}^2} \prod_{j=1}^{km} p(i_j - i_{j-1}, z_{s(j-1)} + \bar{x}_{c(j-1)}, z_{s(j)} + \bar{x}_{c(j)}) \end{aligned} \quad (8.1)$$

where  $s$  runs over the set of maps  $S(k, m)$  from  $\{1, \dots, km\}$  to  $\{1, \dots, m\}$  such that  $s^{-1}(j) = k$  for each  $1 \leq j \leq m$ ,  $c(j) = \sum_{i=1}^j 1_{\{s(i)=s(j)\}}$ ,  $\bar{x}_j = \sum_{l=2}^j x_l$  and  $\bar{x}_1 = 0$ . Here we use the conventions  $i_0 = 0, z_0 = 0, c(0) = 0$ . Setting

$$g_n(x) = \sum_{i=0}^n p(i, 0, x) \leq c \log n \quad (8.2)$$

and using the obvious fact that  $\sum_{x \in \mathbb{Z}^2} g_n(x) = \sum_{x \in \mathbb{Z}^2} \sum_{i=0}^n p(i, 0, x) = n+1$  we can bound (8.1) by

$$\begin{aligned} & \sum_{s \in S(k, m)} \sum_{z_1, \dots, z_m \in \mathbb{Z}^2} \prod_{j=1}^{km} g_n(z_{s(j)} + \bar{x}_{c(j)} - z_{s(j-1)} - \bar{x}_{c(j-1)}) \\ & \leq c(\log n)^{(k-1)m} \sum_{s \in S} \sum_{z_1, \dots, z_k \in \mathbb{Z}^2} \prod_{j: c(j)=1} g_n(z_{s(j)} + \bar{x}_{c(j)} - z_{s(j-1)} - \bar{x}_{c(j-1)}) \\ & \leq cn^m (\log n)^{(k-1)m}. \end{aligned} \quad (8.3)$$

□

**Proof of Lemma 7.3.** Using Proposition 2.1 it suffices to prove (7.4) for all  $k$ .

Let

$$C_k(n, x) = [B_k(n, x) - B_k(n-1, x)] = \sum_{0 \leq i_1 < i_2 < \dots < i_k = n} \prod_{j=2}^k 1_{(X_{i_j} = X_{i_{j-1}} + x_j)}.$$

If we show

$$\mathbb{E} C_k(i, x)^p \leq c(\log n)^{(k-1)p}, \quad i \leq n, \quad (8.4)$$

then using

$$Y_k(n)^p \leq \sum_{i=1}^n c(p) C_k(i, x)^p,$$

we are done.

But

$$\begin{aligned}
& \mathbb{E} \left\{ \left( \sum_{0 \leq i_1 < i_2 < \dots < i_k = n} \prod_{j=2}^k 1_{(X_{i_j} = X_{i_{j-1}} + x_j)} \right)^m \right\} \\
&= \sum_{s \in S(k-1, m)} \sum_{(i_1, \dots, i_{(k-1)m}) \in D(s)} \sum_{z_1, \dots, z_m, y \in \mathbb{Z}^2} \\
& \quad \prod_{j=1}^{(k-1)m} p(i_j - i_{j-1}, z_{s(j-1)} + \bar{x}_{c(j-1)}, z_{s(j)} + \bar{x}_{c(j)}) \\
& \quad \left( \prod_{j: c(j)=k-1} 1_{(y = z_{s(j)} + \bar{x}_k)} \right) p(n - i_{(k-1)m} + \bar{x}_{k-1}, z_{s((k-1)m)}, y)
\end{aligned} \tag{8.5}$$

where  $S(k-1, m)$ ,  $c(j)$ ,  $\bar{x}_j$  are defined in the last section and for each  $s \in S(k-1, m)$

$$\begin{aligned}
D(s) = \{ & (i_1, \dots, i_{(k-1)m}) : 0 \leq i_1 \leq \dots \leq i_{(k-1)m} < n \text{ and} \\
& i_{j-1} < i_j \text{ whenever } s(j-1) = s(j) \}.
\end{aligned}$$

We can then see that (8.5) equals

$$\begin{aligned}
& \sum_{s \in S(k-1, m)} \sum_{(i_1, \dots, i_{(k-1)m}) \in D(s)} \sum_{y \in \mathbb{Z}^2} \\
& \quad p(i_1, 0, y - \bar{x}_k) \prod_{j=2}^{(k-1)m} p(i_j - i_{j-1}, \bar{x}_{c(j-1)}, \bar{x}_{c(j)}) p(n - i_{(k-1)m}, 0, x_k) \\
& \leq c(\log n)^{(k-1)m}
\end{aligned}$$

which is (7.4). □

## 9. Proof of Lemma 7.4.

This proof is substantially different from the proof of Lemma 4.5.

**Proof of Lemma 7.4.** We use induction on  $k$ . We already know (7.6) for  $k = 2$ . Thus assume (7.6) has been proved with  $k$  replaced by  $i$  for all  $i \leq k-1$ . Then as explained above in the proof of Proposition 7.1 we will have that (7.10) holds with  $k$  replaced by  $i$  for all  $i \leq k-1$ .

We will show that

$$\mathbb{E} [ |\tilde{U}_{k,n}(m, x) - \tilde{U}_{k,n}(m, x)|^p ] \leq c(\log n)^{(k-1)p} n^{p+k-1} \left( \frac{|x - x'|}{\sqrt{n}} \right)^{8^{-k} 2p} \tag{9.1}$$

for  $m \leq n$ . This and the inequality

$$\mathbb{E} [\max_{m \leq n} |C_m|^p] \leq \mathbb{E} \sum_{m=1}^n |C_m|^p = \sum_{m=1}^n \mathbb{E} [|C_m|^p] \leq n \max_{m \leq n} \mathbb{E} [|C_m|^p]$$

yields (7.6).

Abbreviating  $\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) := \tilde{B}_{k-1,n}(i, x_{k^c}) - \tilde{B}_{k-1,n}(i-1, x_{k^c})$ , where  $x_{k^c}$  is the same as  $(x_2, \dots, x_{k-1})$ , we have

$$\begin{aligned} & \|\tilde{U}_{k,n}(m, x) - \tilde{U}_{k,n}(m, x')\|_p \\ & \leq \left\| \sum_{i=1}^m \left( G(X_m - X_i - x_k) - G(X_m - X_i - x'_k) \right) \left( \Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) \right) \right\|_p \\ & \quad + \left\| \sum_{i=1}^m \left( G(X_m - X_i - x'_k) - G(e_1 \sqrt{n}) \right) \left( \Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c}) \right) \right\|_p. \end{aligned} \quad (9.2)$$

Then with  $m \leq n$

$$\begin{aligned} & \left\| \sum_{i=1}^m \left( G(X_m - X_i - x_k) - G(X_m - X_i - x'_k) \right) \left( \Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) \right) \right\|_p \\ & \leq \left\| \sum_{i=1}^m |G(X_m - X_i - x_k) - G(X_m - X_i - x'_k)| \right\|_{2p} \|\tilde{Y}_{k-1}(m)\|_{2p} \\ & \leq cn \left| \frac{x - x'}{\sqrt{n}} \right|^{2/3} n^{1/2p} (\log n)^{(k-2)}. \end{aligned} \quad (9.3)$$

by (5.9) and (7.5).

After interchanging  $x'$  and  $x$  for convenience it remains to bound

$$\begin{aligned} & \left\| \sum_{i=1}^m \left( G(X_m - X_i - x_k) - G(e_1 \sqrt{n}) \right) \left( \Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c}) \right) \right\|_p \\ & \leq \left\| \sum_{i=1}^m G(X_m - X_i - x_k) \left( \Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c}) \right) \right\|_p \\ & \quad + \left\| G(\sqrt{n}e_1) [\tilde{B}_{k-1,n}(m, x_{k^c}) - \tilde{B}_{k-1,n}(m, x'_{k^c})] \right\|_p \end{aligned} \quad (9.4)$$

Using Proposition 2.1 and our inductive hypothesis concerning (7.10) we see that

$$\begin{aligned} & \left\| G(\sqrt{n}e_1) [\tilde{B}_{k-1,n}(m, x_{k^c}) - \tilde{B}_{k-1,n}(m, x'_{k^c})] \right\|_p \\ & \leq c(\log n)^{(k-1)} n^{1+(k-1)/p} \left( \frac{|x - x'|}{\sqrt{n}} \right)^{8^{-k+1}}. \end{aligned} \quad (9.5)$$

To complete the proof of (9.1) it therefore suffices to show that

$$\begin{aligned} & \left\| \sum_{i=1}^m G(X_m - X_i - x) [\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c})] \right\|_p \\ & \leq c(\log n)^{(k-1)} n^{1+(k-1)/p} \left( \frac{|x - x'|}{\sqrt{n}} \right)^{8^{-k} 2}. \end{aligned} \quad (9.6)$$

By Propositions 2.1 and 2.3,  $G(x)$  is bounded above (but  $G(x) \rightarrow -\infty$  as  $|x| \rightarrow \infty$ ).

Let

$$J(x) = G(x) \vee (-9 \log m), \quad H(x) = G(x) - J(x).$$

Let  $K_i = J(X_m - X_i - x)$  for  $i = 0, \dots, m$  and let  $K_i = J(X_m - x)$  for  $i < 0$ . Let  $B$  be a small positive real to be chosen later and let

$$L_i = \frac{K_i + \dots + K_{i-Bm}}{Bm}.$$

We see that

$$\begin{aligned} & \left| \sum_{i=1}^m (K_i - L_i) [\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c})] \right| \\ & \leq \tilde{Y}_{k-1}(n) \sum_{i=1}^n |K_i - L_i|. \end{aligned} \quad (9.7)$$

Since  $J$  is bounded in absolute value by  $c \log m$ , the same is true for  $K_i$  and  $L_i$  for any  $i$ , i.e.

$$\sup_i |K_i - L_i| \leq c \log m. \quad (9.8)$$

Note that  $L_i$  and  $K_i$  are independent of  $\mathcal{F}_h$  for  $i \geq h + Bm$ , and thus

$$\mathbb{E} \left[ \sum_{i=h}^m |K_i - L_i| \mid \mathcal{F}_h \right] \leq \mathbb{E} \sum_{i=h+2Bm}^m |K_i - L_i| + cBm \log m. \quad (9.9)$$

Now by Proposition 2.2

$$\begin{aligned} & \mathbb{E} |J(X_m - X_i - x) - J(X_m - X_j - x)| \\ & \leq \mathbb{E} |G(X_m - X_i - x) - G(X_m - X_j - x)| \\ & \leq \mathbb{E} \left[ \frac{c|X_i - X_j|^{2/3}}{(1 + |X_m - X_i - x|^2)^{1/3}} + \frac{c|X_i - X_j|^{2/3}}{(1 + |X_m - X_j - x|^2)^{1/3}} \right]. \end{aligned}$$

By (5.6) and symmetry

$$\mathbb{E} \left( (1 + |X_m - X_i - x|^2)^{-1/2} \right) \leq \mathbb{E} \left( (1 + |X_m - X_i|^2)^{-1/2} \right) \leq 1 \wedge c(m-i)^{-1/2}.$$

Then using Holder's inequality in the form  $|\mathbb{E}(fg)| \leq \|f\|_3 \|g\|_{3/2}$  we obtain from the last two displays that

$$\begin{aligned} \mathbb{E} |J(X_m - X_i - x) - J(X_m - X_j - x)| & \quad (9.10) \\ & \leq \frac{c|i-j|^{1/3}}{1 \vee |m-i|^{1/3}} + \frac{c|i-j|^{1/3}}{1 \vee |m-j|^{1/3}}. \end{aligned}$$

Thus for  $i \geq 2Bm$ , summing over  $j$  from  $i - Bm$  to  $i$  and dividing by  $Bm$  shows

$$\mathbb{E} |K_i - L_i| \leq c(Bm)^{1/3} (1 \vee |m-i|)^{-1/3}.$$

Therefore,

$$\mathbb{E} \sum_{i=h+2Bm}^m |K_i - L_i| \leq \sum_{i=h+2Bm}^m \frac{c(Bm)^{1/3}}{(1 \vee |m-i|)^{1/3}} \leq cmB^{1/3}.$$

Recalling (9.8)–(9.9) and then using Proposition 3.1 we have that

$$\mathbb{E} \left| \sum_{i=1}^m |K_i - L_i| \right|^p \leq c(\log m)^p + cm^p B^{p/3} \leq c(\log n)^p n^p B^{p/3} \quad (9.11)$$

for  $n$  large. Combining with this with (9.7), (7.5) and Cauchy-Schwarz, the left hand side of (9.7) is bounded in  $L^p$  norm by

$$c(\log n)^{k-1} n^{1+(1/2p)} B^{1/3}. \quad (9.12)$$

We use summation by parts on

$$\sum_{i=1}^m L_i [\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c})] \quad (9.13)$$

and we see that it is equal to

$$\begin{aligned} & L_m [\tilde{B}_{k-1,n}(m, x_{k^c}) - \tilde{B}_{k-1,n}(m, x'_{k^c})] \\ & - \sum_{i=1}^m [\tilde{B}_{k-1,n}(i-1, x_{k^c}) - \tilde{B}_{k-1,n}(i-1, x'_{k^c})] [L_i - L_{i-1}]. \end{aligned} \quad (9.14)$$

Write  $w = |x_{k^c} - x'_{k^c}|/\sqrt{n} \leq 1$ . Using the fact that  $L_m$  is bounded by  $c \log m$  and our inductive hypothesis concerning (7.10), we can bound the  $L^p$  norm of the first term of (9.14) by

$$c(\log n)^{k-1} n^{1+(k-1)/p} w^{8^{-k+1}}.$$



Since  $K_i$  is bounded by  $c \log n$ , then  $L_i - L_{i-1}$  is bounded by  $c \log n / (Bn)$ . Hence using once again our inductive hypothesis concerning (7.10)

$$\begin{aligned} & \| [\tilde{B}_{k-1,n}(i-1, x_{k^c}) - \tilde{B}_{k-1,n}(i-1, x'_{k^c})] [L_i - L_{i-1}] \|_p \\ & \leq \frac{c \log n}{Bn} \| \tilde{B}_{k-1,n}(i-1, x_{k^c}) - \tilde{B}_{k-1,n}(i-1, x'_{k^c}) \|_p \\ & \leq \frac{c \log n}{Bn} (\log n)^{k-2} n^{1+(k-1)/p} w^{8^{-k+1}}. \end{aligned}$$

Since there are  $n$  summands in the sum in (9.14), we bound the  $L^p$  norm of the left hand side of (9.13) by

$$\frac{c}{B} (\log n)^{k-1} n^{1+(k-1)/p} w^{8^{-k+1}}. \quad (9.15)$$

Notice that

$$\left| \sum_{i=1}^m H(X_m - X_i - x) [\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c})] \right| \leq m \tilde{Y}_{k-1}(n) \sup_{1 \leq i \leq m} |H(X_m - X_i - x)|. \quad (9.16)$$

By Proposition 2.3,  $H(z)$  is 0 unless  $|z| \geq e^{8 \log m}$ . By hypothesis we have  $|x| \leq \sqrt{n}$ . Therefore using  $|H(z)|^{2p} \leq c |\log z|^{2p} \leq c |z|$  for  $|z| \geq e^{8 \log m}$

$$\begin{aligned} \mathbb{E} |H(X_m - X_i - x)|^{2p} & \leq c(p) \mathbb{E} [|X_m - X_i - x|; |X_m - X_i - x| \geq e^{8 \log m}] \\ & \leq c(p) e^{-8 \log m} \mathbb{E} |X_m - X_i - x|^2 \\ & \leq c(p) m^2 e^{-8 \log m} = c(p) m^{-6}. \end{aligned}$$

Hence  $\mathbb{E} \sup_{i \leq m} |H(X_m - X_i - x)|^{2p} \leq c(p) m^{-5}$ . Since  $w \geq 1/\sqrt{n}$ , then this estimate, (9.16), (7.5) and Cauchy-Schwarz imply that the left hand side of (9.16) is bounded in  $L^p$  norm by  $c(\log n)^{k-2} n^{1/2p} \leq c(\log n)^{k-2} n^{1+1/2p} w^2 \leq c(\log n)^{k-1} n^{1+1/2p} w^{8^{-k+1}}$ .

Combining our estimates (9.12), (9.15), and our last estimate for (9.16), we have

$$\begin{aligned} & \left\| \sum_{i=1}^m G(X_m - X_i - x) [\Delta_i \tilde{B}_{k-1,n}(i, x_{k^c}) - \Delta_i \tilde{B}_{k-1,n}(i, x'_{k^c})] \right\|_p \\ & \leq c(\log n)^{(k-1)} n^{1+(k-1)/p} [w^{8^{-k+1}} + B^{1/3} + w^{8^{-k+1}} B^{-1}]. \end{aligned} \quad (9.17)$$

If we take  $B = w^{6(8^{-k})}$ , we obtain (9.6). Together with (9.2)–(9.5) we obtain (9.1).  $\square$

## 10. Other results.

A.  $L^2$  norms. By Section 3 of [5] we see that we can choose  $W_t$  and  $X_n$  such that

$$\left\| \sup_{s \leq 1} |X_s^n - W_s^n| \right\|_2 = o(n^{-\zeta})$$

for some  $\zeta > 0$ . If we then use this (in place of (1.7)), our proof shows that we obtain

$$\|\tilde{\beta}_k(n, 0) - \tilde{\gamma}_k(1, 0, n)\|_2 = o(n^{-\eta}) \quad (10.1)$$

for some  $\eta > 0$ .

*B. A correction.* We take this opportunity to correct an error in [3]. In the statement of (8.3) in Theorem 8.1 of that paper,  $G^\vee := \max_{1 \leq j \leq k-1} |G(x_j)|$  should be replaced by  $N^\vee := \max_{1 \leq j \leq k-1} |x_j|^{-1}$ . The term  $G^\vee$  also needs to be replaced by  $N^\vee$  throughout the proof of (8.3).

Proposition 9.2 of that paper is correct as stated. Where the proof of this proposition says to follow the lines of the proof of (8.3), it is to be understood that here one uses  $G^\vee$  throughout.

For the convenience of the interested reader we give a complete proof of that proposition in the following Appendix.

### Appendix. Proof of Proposition 9.2 of [3].

The proof of Proposition 9.2 in [3] is perhaps a bit confusing due to an error in the statement of (8.3) in Theorem 8.1 of that paper. This Appendix provides a complete proof of Proposition 9.2 of [3].

Write  $g(y) = \frac{1}{\pi} \log(1/|y|)$ ,  $\tilde{\gamma}_1(t, x) = t$ , and for  $x = (x_2, \dots, x_k) = (x_{k^c}, x_k) \in (\mathbb{R}^2)^{k-1}$ , set

$$g^\vee(x) = \max_{2 \leq i \leq k} |g(x_i)|$$

and

$$\bar{U}_k(t, x) = \int_0^t g(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}).$$

**Proposition A.1.** *Let  $M \geq 1$ , let  $x, x' \in (\mathbb{R}^2)^{k-1}$  with  $|x_i|, |x'_i| \leq M$  for  $i = 2, \dots, k$ , and let  $g^+ = g^\vee(x) + g^\vee(x') + 1$ . There exist  $a_k$  and  $\nu_k$  such that for  $k \geq 2$*

$$(a) \quad \mathbb{E} |\bar{U}_k(t, x) - \bar{U}_k(t, x')|^p \leq c(g^+)^{\nu_k p} |x - x'|^{a_k p}.$$

$$(b) \quad \mathbb{E} |\bar{U}_k(t, x) - \bar{U}_k(s, x)|^p \leq c(g^+)^{\nu_k p} |t - s|^{a_k p}.$$

Except for the restriction on the size of  $x, x'$ , this is Proposition 9.2 of [3] translated to the notation of this paper. Using the argument of [3] this is sufficient to prove the joint continuity of  $\tilde{\gamma}_k(t, x)$  over  $t \in [0, 1], x \in (B(0, M))^{k-1}$  for each  $k$  and  $M$ . For almost every path of Brownian motion,  $\{W_s : s \in [0, 1]\}$  is contained in  $B(0, M)$  for some  $M$  (depending on the path), and hence for  $|x| > M$  we have  $\tilde{\gamma}_k(t, x) = 0$ . The joint continuity of  $\tilde{\gamma}_k(t, x)$

over  $t \in [0, 1], x \in (\mathbb{R}^2)^{k-1}$  follows. For the purposes of this paper we only need the case  $M = 1$ .

Note that renormalization allows us to use  $\bar{U}_k(t, x)$  in place of

$$U_k^*(t, x) = \int_0^t [g(W_t - W_r - x_k) - g(-x_k)] \tilde{\gamma}_{k-1}(dr, x_{k^c}).$$

If one were to try to use  $U_k^*(t, x)$  in Proposition A.1, the right hand sides of (a) and (b) would have to have  $g^+$  replaced by  $N^\vee$ , which is not a good enough bound for the joint continuity argument.

**Proof.** Since  $g^+$  is infinite if any component of  $x$  or  $x'$  is zero, we may assume that no component of either is 0.

Let  $A \in (0, \frac{1}{2}]$  be chosen later and let

$$g_A(x) = (g(x) \wedge \frac{1}{\pi} \log(1/A)) \vee (-\frac{1}{\pi} \log(1/A)),$$

$$h_A(x) = [g(x) - g_A(x)]1_{(|x| < A)}, \quad j_A(x) = [g_A(x) - g(x)]1_{(|x| > A^{-1})},$$

so  $g = g_A + h_A - j_A$ . With  $C_k = C \cup \{k\}$  set

$$L_k(t, x) = \sum_{C \subset \{2, \dots, k-1\}} \left( \prod_{i \in C} |g(x_i)| \right) \gamma_{k-1-|C|}(t, x_{C_k^c}).$$

The proof is by induction. We start with  $k = 2$ . In preparation for general  $k$  we retain the general notation, but note that when  $k = 2$ , we have  $L_2(t, x) = t$ ,  $x_{k^c}$  is superfluous, and we have  $\tilde{\gamma}_{k-1}(dr, x_{k^c}) = dr$ .

$$\begin{aligned} & \bar{U}_k(t, x) - \bar{U}_k(t, x') \\ &= \int_0^t [g_A(W_t - W_r - x_k) - g_A(W_t - W_r - x'_k)] \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\ & \quad + \int_0^t h_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\ & \quad - \int_0^t h_A(W_t - W_r - x'_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\ & \quad - \int_0^t j_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\ & \quad + \int_0^t j_A(W_t - W_r - x'_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\ &=: I_1 + I_2 - I_3 - I_4 + I_5. \end{aligned} \tag{A.1}$$

If we connect  $x, x'$  by a curve  $\Gamma$  of length  $c|x - x'|$  that never gets closer to 0 than  $|x| \wedge |x'|$ , use the fact that  $|\nabla g_A| \leq A^{-1}$ , and use inequality (8.1) of [3] (this is only needed for  $k > 2$ )

$$\mathbb{E}|I_1|^p \leq cA^{-p}|x - x'|^p \mathbb{E}L_k(t, x)^p \leq cA^{-p}(g^+)^{\nu_1 p}|x - x'|^p.$$

By Proposition 5.2 of [3], for some constants  $b_1$  and  $\nu'_1$

$$\mathbb{E}|I_2|^p \leq cA^{b_1 p}(g^+)^{\nu'_1 p} \quad (\text{A.2})$$

and similarly for  $I_3$ .

We next turn to  $I_4$ . Standard estimates on Brownian motion tells us that

$$\mathbb{P}\left(\sup_{0 \leq r \leq t \leq 1} |W_t - W_r| > \lambda\right) \leq ce^{-c'\lambda^2}. \quad (\text{A.3})$$

Since  $|x_k| \leq M$ , it follows that

$$\mathbb{E}\left[\sup_{0 \leq r \leq t \leq 1} |\log(1/|W_t - W_r - x_k|)|^p\right] \leq c(p, M) \quad (\text{A.4})$$

for each  $p \geq 1$ . If  $|A| \leq (2M)^{-1}$ , then  $|W_t - W_r - x_k| \geq A^{-1}$  only if  $|W_t - W_r| \geq (2A)^{-1}$ . So by Cauchy-Schwarz, (A.3), and (A.4),

$$\begin{aligned} & \mathbb{E}\left[\sup_{0 \leq r \leq t \leq 1} |j_A(W_t - W_r - x_k)|^p\right] \\ & \leq c\mathbb{E}\left[\sup_{0 \leq r \leq t \leq 1} |\log(1/|W_t - W_r - x_k|)|^p \mathbf{1}_{(\sup_{0 \leq r \leq t \leq 1} |W_t - W_r - x_k| \geq A^{-1})}\right] \\ & \leq c(\mathbb{E}\left[\sup_{0 \leq r \leq t \leq 1} |\log(1/|W_t - W_r - x_k|)|^{2p}\right])^{1/2} (\mathbb{P}(\sup_{0 \leq r \leq t \leq 1} |W_t - W_r| \geq (2A)^{-1}))^{1/2} \\ & \leq cA^p \end{aligned}$$

for each  $p \geq 1$ . Using the fact that

$$|I_4| \leq \left(\sup_{0 \leq r \leq t \leq 1} |j_A(W_t - W_r - x_k)|\right)L_k(t, x),$$

another application of Cauchy-Schwarz shows that

$$\mathbb{E}|I_4|^p \leq (g^+)^{\bar{\nu}_1 p} A^p.$$

$I_5$  is handled the same way.

Combining shows the left hand side of (A.1) is bounded by

$$c(g^+)^{\nu_2 p}[A^{-p}|x - x'|^p + A^{b_1 p} + A^p]$$

for some constant  $\nu_2$ , and we obtain (a) for  $k = 2$  by setting  $A = |x - x'|^{1/2} \wedge (2M)^{-1}$ .

Next we look at (b) for the  $k = 2$  case. We write

$$\begin{aligned}
& \bar{U}_k(t, x) - \bar{U}_k(s, x) \\
&= \int_0^s [g_A(W_t - W_r - x_k) - g_A(W_s - W_r - x_k)] \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad + \int_s^t g_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad + \int_0^t h_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad - \int_0^s h_A(W_s - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad - \int_0^s j_A(W_s - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad + \int_0^s j_A(W_s - W_r - x'_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&=: I_6 + I_7 + I_8 - I_9 - I_{10} + I_{11}.
\end{aligned} \tag{A.5}$$

Using Cauchy-Schwarz, for  $s, t \leq 1$  we have for some constant  $\nu_1''$

$$\mathbb{E} |I_6|^p \leq cA^{-p} s^p \left( \mathbb{E} |W_t - W_s|^{2p} \right)^{1/2} \left( \mathbb{E} L(s, x_{k^c})^{2p} \right)^{1/2} \leq cA^{-p} (g^+)^{\nu_1'' p} |t - s|^{p/2}.$$

We bound  $I_7$  by

$$\mathbb{E} |I_7|^p \leq c(\log(1/A))^p |t - s|^p (g^+)^{\nu''' p}.$$

We bound  $I_8$  and  $I_9$  just as we did  $I_2$  and bound  $I_{10}$  and  $I_{11}$  as we did  $I_4$ . Combining, the left hand side of (A.5) is bounded by

$$c(g^+)^{\nu_2 p} [A^{-p} |t - s|^{p/2} + A^{b_1 p} + A^p],$$

and (b) follows by setting  $A = |t - s|^{1/4} \wedge (2M)^{-1}$ .

We now turn to the case when  $k > 2$ . We suppose (a) and (b) hold for  $k - 1$  and prove them for  $k$ . We prove (a) in two cases, when  $x_{k^c} = x'_{k^c}$  and when  $x_k = x'_k$ ; the general case follows by the triangle inequality. Suppose first that  $x_{k^c} = x'_{k^c}$ . Using the induction hypothesis, the proof is almost exactly the same as the proof of (a) in the case  $k = 2$ .

Suppose next that  $x_k = x'_k$ . Let

$$V_A = \{|W_{s+u} - W_u| \geq u^{1/4}/A \text{ for some } s, u \in [0, 1]\}.$$

Standard estimates on Brownian motion show that

$$\mathbb{P}(V_A) \leq c_1 e^{-c_2/A^2}.$$

We write

$$\begin{aligned}
& \bar{U}_k(t, x) - \bar{U}_k(t, x') \\
&= 1_{V_A} \int_0^t g_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad - 1_{V_A} \int_0^t g_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x'_{k^c}) \\
&\quad + \int_0^t h_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad - \int_0^t h_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x'_{k^c}) \\
&\quad - \int_0^t j_A(W_t - W_r - x_k) \tilde{\gamma}_{k-1}(dr, x'_{k^c}) \\
&\quad + \int_0^t j_A(W_t - W_r - x'_k) \tilde{\gamma}_{k-1}(dr, x'_{k^c}) \\
&\quad + 1_{V_A^c} \int_0^t g_A(W_t - W_r - x_k) [\tilde{\gamma}_{k-1}(dr, x_{k^c}) - \tilde{\gamma}_{k-1}(dr, x'_{k^c})] \\
&=: I_{12} - I_{13} + I_{14} - I_{15} - I_{16} + I_{17} + I_{18}.
\end{aligned} \tag{A.6}$$

Since  $|g_A| \leq \log(1/A)$ , for some constant  $\nu_{k-1}$

$$\begin{aligned}
\mathbb{E} |I_{12}|^p &\leq c(\log(1/A))^p \mathbb{E} [L_k(t, x)^p; V_A] \\
&\leq c(\log(1/A))^p \left( \mathbb{E} L_k(t, x)^{2p} \right)^{1/2} \left( \mathbb{P}(V_A) \right)^{1/2} \leq c(g^+)^{\nu_{k-1} p} A^p,
\end{aligned}$$

and similarly for  $I_{13}$ . We bound  $I_{14}$  and  $I_{15}$  just as we did  $I_2$  and bound  $I_{16}$  and  $I_{17}$  as we did  $I_4$ .

We turn to  $I_{18}$ . Let  $f(r) = g_A(W_t - W_r - x_k)$  and

$$f_A(t) = \frac{1}{A^{12}} \int_t^{t+A^{12}} f(u) du.$$

On  $V_A^c$  we have

$$\begin{aligned}
|f(r) - f(s)| &= |g_A(W_t - W_r - x_k) - g_A(W_t - W_s - x_k)| \\
&\leq A^{-1} |W_r - W_s| \leq A^{-2} |r - s|^{1/4},
\end{aligned}$$

and therefore

$$|f(t) - f_A(t)| \leq A^{-2} (A^{12})^{1/4} = A.$$

Using integration by parts, we write

$$\begin{aligned}
I_{18} &= 1_{V_A^c} \int_0^t [f(r) - f_A(r)] \tilde{\gamma}_{k-1}(dr, x_{k^c}) \\
&\quad - 1_{V_A^c} \int_0^t [f(r) - f_A(r)] \tilde{\gamma}_{k-1}(dr, x'_{k^c}) \\
&\quad + 1_{V_A^c} f_A(t) [\tilde{\gamma}_{k-1}(t, x_{k^c}) - \tilde{\gamma}_{k-1}(t, x'_{k^c})] \\
&\quad - 1_{V_A^c} \int_0^t [\tilde{\gamma}_{k-1}(r, x_{k^c}) - \tilde{\gamma}_{k-1}(r, x'_{k^c})] f_A(dr) \\
&=: I_{19} - I_{20} + I_{21} - I_{22}.
\end{aligned}$$

We bound

$$\mathbb{E} |I_{19}|^p \leq cA^p (g^+)^{\nu'_{k-1} p}$$

for some constant  $\nu'_{k-1}$  and similarly for  $I_{20}$ . By the induction hypothesis and the fact that  $|f_A|$  is bounded by  $\log(1/A) \leq cA^{-p}$ ,

$$\mathbb{E} |I_{21}|^p \leq cA^{-p} (g^+)^{\nu''_{k-1} p} |x_{k^c} - x'_{k^c}|^{a_{k-1} p}.$$

Finally, since  $|f'_A| \leq \|f_A\|_\infty A^{-12} \leq cA^{-13}$ ,

$$\mathbb{E} |I_{22}|^p \leq c(g^+)^{\nu'''_{k-1} p} A^{-13p} |x_{k^c} - x'_{k^c}|^{a'_{k-1} p}.$$

If we combine all the terms, we see that the left hand side of (A.6) is bounded by

$$c(g^+)^{\nu_{k^c} p} [A^{b_k p} + A^{-b'_k p} |x - x'|^{a''_{k-1} p} + A^p].$$

Setting  $A = |x - x'|^{a''_{k-1}/(2b'_k)} \wedge (2M)^{-1}$  completes the proof of (b) for  $k > 2$ .

The proof of (b) for  $k > 2$  is almost identical to the  $k = 2$  case.  $\square$

## References.

- [1] R.F. Bass, *Probabilistic Techniques in Analysis*. Springer, New York, 1995.
- [2] R.F. Bass and D. Khoshnevisan, Local times on curves and uniform invariance principles. *Prob. Th. and rel. Fields* **92** (1992) 465–492.
- [3] R.F. Bass and D. Khoshnevisan, Intersection local times and Tanaka formulas. *Ann. Inst. H. Poincaré* **29** (1993) 419–451.

- [4] R.F. Bass and D. Khoshnevisan, Strong approximations to Brownian local time, *Seminar on Stochastic Processes, 1992*, 43–66, Birkhäuser, Boston, 1993.
- [5] R.F. Bass and J. Rosen, An almost sure invariance principle for the range of planar random walks, *Ann. Probab.*, to appear.
- [6] I. Berkes and W. Philipp, Approximation theorems for independent and weakly dependent random vectors, *Ann. Probab.* **7** (1979) 29–54.
- [7] E. B. Dynkin, Self-intersection gauge for random walks and for Brownian motion. *Ann. Probab.* **16** (1988) 1–57.
- [8] E. B. Dynkin, Regularized self-intersection local times of planar Brownian motion. *Ann. Probab.* **16** (1988) 58–74.
- [9] J.-F. Le Gall, Wiener sausage and self intersection local times. *J. Funct. Anal.* **88** (1990) 299–341.
- [10] J.-F. Le Gall. Propriétés d’intersection des marches aléatoires, I. Convergence vers le temps local d’intersection, *Comm. Math. Phys.* **104** (1986) 471–507.
- [11] J.-F. Le Gall. Some properties of planar Brownian motion, *École d’été de probabilités de St. Flour XX, 1990* Springer-Verlag, Berlin, 1992.
- [12] G. Lawler, *Intersections of random walks*. Birkhäuser, Boston, 1991.
- [13] G. Lawler, in preparation.
- [14] M. Marcus and J. Rosen, *Renormalized self-intersection local times and Wick power chaos processes*, Memoirs of the AMS, Volume 142, Number 675, Providence, AMS, 1999.
- [15] V.V. Petrov, *Sums of Independent Random Variables*. Springer-Verlag, Berlin, 1975.
- [16] J. Rosen, Random walks and intersection local time. *Ann. Probab.* **18** (1990) 959–977.
- [17] J. Rosen, Joint continuity of renormalized intersection local times, *Ann. Inst. H. Poincaré* **32** (1996) 671–700.
- [18] J. Rosen. Dirichlet processes and an intrinsic characterization for renormalized intersection local times, *Ann. Inst. H. Poincaré* **37** (2001) 403–420.
- [19] F. Spitzer, *Principles of Random Walk, 2nd ed.*, Springer, New York, 1976.
- [20] S. R. S. Varadhan. Appendix to Euclidean quantum field theory by K. Symanzyk. In R. Jost, editor, *Local Quantum Theory*. Academic Press, Reading, MA, 1969.



R. Bass Address: Department of Mathematics, University of Connecticut,  
Storrs, CT 06269-3009 (bass@math.uconn.edu)

J. Rosen Address: Department of Mathematics, College of Staten Island, CUNY,  
Staten Island, NY, 10314 (jrosen3@earthlink.net)