


- (1) (15 points) A hot air balloon rises vertically upwards from a distance of two miles away. When you see the balloon at an angle of $\pi/6$ radians, the angle is changing at a rate of 0.1 radians/s. How fast is the balloon rising?




$$\tan \theta = \frac{h}{2}$$

$$\sec^2 \theta \frac{d\theta}{dt} = \frac{1}{2} \frac{dh}{dt}$$

$$\frac{dh}{dt} = 2 \sec^2\left(\frac{\pi}{6}\right) \cdot 0.1 = \frac{0.2}{\left(\frac{\sqrt{3}}{2}\right)^2} \text{ miles/s}$$

$$= \frac{0.8}{3} \approx 0.2667$$

(2) (25 points) Consider the function $f(x)$ defined by the following graph.



- 4(a) Label all regions where $f(x) < 0$.
 4(b) Label all regions where $f'(x) > 0$.
 4(c) What is $\lim_{x \rightarrow -\infty} f(x)$?
 4(d) What is $\lim_{x \rightarrow \infty} f'(x)$?
 5(e) Sketch a graph of $f'(x)$ on the figure.
 4(f) Label the approximate locations of all points of inflection.

c) approx -1

d) 0

- (3) (15 points) The value of $\tan x$ at $\pi/4$ is 1. Use a linear approximation to estimate $\tan(0.8)$. Do you consider this to be a good approximation?

$$\Delta f \approx f'(\pi/4) (0.8 - \pi/4) \quad \Delta x = 0.8 - \pi/4 \approx 0.0146$$

$$f(x) = \tan(x)$$

$$f'(x) = \sec^2(x) \quad f'(\pi/4) = 2$$

$$\text{so } \Delta f \approx f'(\pi/4) \Delta x = 0.0292$$

$$\text{so } \tan(0.8) \approx 1 + 0.0292 = 1.0292$$

$$\text{actual value: } \tan(0.8) = 1.02964\dots$$

$$\text{good approximation: error is } |1.0292 - 1.02964| \approx 0.0004$$

$$\text{percentage error is } \frac{100 \times 0.0004}{1.02964\dots} \approx 0.04\%$$

(4) (25 points) Consider the function

$$f(x) = \frac{1}{x^2 - 3x + 2}$$

- Find all vertical and horizontal asymptotes of the function.
- Find all critical points of the function.
- Determine the intervals where $f(x)$ is increasing and decreasing.
- Use the 2nd derivative test to attempt to identify all local maxima and minima.
- Sketch the function and label all relative maxima and minima.

a) vertical asymptotes: $x^2 - 3x + 2 = (x-2)(x-1)$

so vertical asymptotes at $x=1, 2$


horizontal asymptotes: $\lim_{x \rightarrow \infty} f(x) = 0$ $\lim_{x \rightarrow -\infty} f(x) = 0$

b) $f'(x) = \frac{-(2x-3)}{(x^2-3x+2)^2}$ $f'(x) = 0$ $x = \frac{3}{2}$


c) $f'(x) > 0, x < \frac{3}{2}$ increasing
 $f'(x) < 0, x > \frac{3}{2}$ decreasing

d) $f''(x) = \frac{(x^2-3x+2)^2(-2) - (2(x^2-3x+2)(2x-3))(-2x+3)}{(x^2-3x+2)^4}$

$f''(\frac{3}{2}) = \frac{(+)(-) - (0)}{(+)} < 0$ so local max



- (5) (15 points) A cylindrical can of volume 1ft^3 is to be constructed, where the material for the top and bottom costs three times as much as the material for the sides. Find the dimensions which minimize the cost of the can.



$$\text{volume: } 1 = \pi r^2 h \Rightarrow h = \frac{1}{\pi r^2}$$

$$\text{cost: } C = 3(2\pi r^2) + 2\pi r h$$

$$C = 6\pi r^2 + \frac{2\pi r}{\pi r^2} = 6\pi r^2 + \frac{2}{r}$$

$$\frac{dC}{dr} = 12\pi r - \frac{2}{r^2} = 0$$

$$r^3 = \frac{1}{6\pi} \quad r = \sqrt[3]{\frac{1}{6\pi}} \approx 0.375751$$

$$h \approx 2.4545$$

(6) (25 points) Compute the following limits. Show all work.

(a) $\lim_{x \rightarrow -\infty} \frac{x^4 + 3x^3 - 3}{(1 - 2x^2)^2}$

(b) $\lim_{x \rightarrow \infty} \frac{2x}{\sqrt{5x + x^2}}$

(c) $\lim_{x \rightarrow 0} \frac{e^{3x} - 1}{\sin x}$

(d) $\lim_{x \rightarrow 0} \frac{1}{1 - \cos x} - \frac{1}{x^2}$

(e) $\lim_{x \rightarrow 0} x^{\sin x}$

$$a) = \lim_{x \rightarrow -\infty} \frac{1 + 3/x - 3/x^4}{(\frac{1}{x^2} - 2)^2} = \frac{1}{4}$$

$$b) = \lim_{x \rightarrow \infty} \frac{2}{\sqrt{5/x + 1}} = 2$$

$$c) = \lim_{x \rightarrow 0} \frac{3e^{3x}}{\cos x} = 3.$$

(L'Hopital)

$$d) = \lim_{x \rightarrow 0} \frac{x^2 - 1 + \cos x}{(1 - \cos x)x^2} \stackrel{\text{(L'Hopital)}}{=} \lim_{x \rightarrow 0} \frac{2x - \sin x}{2x - 2x \cos x + x^2 \sin x}$$

$$\stackrel{\text{L'Hopital}}{=} \lim_{x \rightarrow 0} \frac{2 - \cos x}{2 - 2\cos x + 2x \sin x + 2x \sin x + x^2 \cos x} = \frac{1}{2 - 2 = 0} \rightarrow \neq \infty.$$

$$e) = \lim_{x \rightarrow 0} e^{\ln x \cdot \sin x} \quad \lim_{x \rightarrow 0} \ln x \cdot \sin x = \lim_{x \rightarrow 0} \frac{\ln x}{1/\sin x}$$

$$= \lim_{x \rightarrow 0} \frac{1/x}{\frac{+\cos x}{\sin^2 x}} = \lim_{x \rightarrow 0} \frac{\sin x}{x} \tan x = 0. \text{ so } \lim_{x \rightarrow 0} e^{\ln x \cdot \sin x} = 1.$$