

**SEMI-FREDHOLM SOLVABILITY IN THE FRAMEWORK
OF SINGULAR SOLUTIONS FOR THE (3+1)-D
PROTTER-MORAWETZ PROBLEM**

NEDYU POPIVANOV¹, TODOR POPOV¹ AND ALLEN TESDALL²

¹Department of Mathematics and Informatics
University of Sofia, 1164 Sofia, Bulgaria
e-mails: nedyu@fmi.uni-sofia.bg, topover@fmi.uni-sofia.bg

²Department of Mathematics, City University of New York
College of Staten Island, Staten Island, NY 10314, USA
e-mail: allen.tesdall@csi.cuny.edu

ABSTRACT. For the four-dimensional nonhomogeneous wave equation boundary value problems that are multi-dimensional analogues of Darboux problems in the plane are studied. It is known that for smooth right-hand side functions the unique generalized solution may have a strong power-type singularity at only one point. This singularity is isolated at the vertex O of the boundary light characteristic cone and does not propagate along the bicharacteristics. The present paper describes asymptotic expansions of the generalized solutions in negative powers of the distance to O . Some necessary and sufficient conditions for existence of bounded solutions are proven and additionally a priori estimates for the singular solutions are obtained.

1. INTRODUCTION

In the present paper, boundary value problems for the wave equation in \mathbb{R}^4

$$(1.1) \quad u_{x_1x_1} + u_{x_2x_2} + u_{x_3x_3} - u_{tt} = f(x, t)$$

with points $(x, t) = (x_1, x_2, x_3, t)$ are studied in the domain

$$\Omega = \{(x, t) : 0 < t < 1/2, t < \sqrt{x_1^2 + x_2^2 + x_3^2} < 1 - t\},$$

bounded by the two characteristic cones

$$\Sigma_1 = \{(x, t) : 0 < t < 1/2, \sqrt{x_1^2 + x_2^2 + x_3^2} = 1 - t\},$$

$$\Sigma_2 = \{(x, t) : 0 < t < 1/2, \sqrt{x_1^2 + x_2^2 + x_3^2} = t\}$$

and the ball $\Sigma_0 = \{t = 0, \sqrt{x_1^2 + x_2^2 + x_3^2} < 1\}$, centered at the origin $O : x = 0, t = 0$. The following BVPs were proposed by M. Protter [33]:

2000 Mathematics Subject Classification. 35L05, 35L20, 35D05, 35D10, 35C10, 35B40 .

Key words and phrases. Wave equation, boundary value problems, generalized solution, semi-Fredholm solvability, asymptotic expansion.

Problem P2. Find a solution of the wave equation (1.1) in Ω which satisfies the boundary conditions

$$P2 : \quad u_t|_{\Sigma_0} = 0, \quad u|_{\Sigma_1} = 0;$$

and its the adjoint problem:

Problem P2*. Find a solution of the wave equation (1.1) in Ω which satisfies the adjoint boundary conditions

$$P2^* : \quad u_t|_{\Sigma_0} = 0, \quad u|_{\Sigma_2} = 0.$$

Protter [33] formulated in 1952 some versions of $P2$ and $P2^*$ in \mathbb{R}^3 (i.e., in (2+1)-D case) as a multi-dimensional analogue of the planar Darboux problems with boundary data prescribed on one characteristic and on the noncharacteristic segment. Initially the expectation was that such BVPs are classical solvable for very smooth right-hand side functions. However, soon it became clear that contrary to this traditional belief, unlike the plane Darboux problem, Protter's problems are not well posed. The reason is that the homogeneous adjoint Problem $P2^*$ has an infinite number of nontrivial classical solutions (Tong Kwang-Chang [36], Popivanov and Schneider [26], Khe Kan Cher [19]). It is known from [27] that for each $n \in \mathbb{N}$ there exists a right-hand side function $f \in C^n(\overline{\Omega})$ of the wave equation, for which the uniquely determined generalized solution of Problem $P2$ has a strong power-type singularity like r^{-n} at the origin O .

In the present paper we examine the exact behavior of the singular solutions of Problem $P2$. In the case when the right-hand side function f is harmonic polynomial, the Problem $P2$ is Fredholm and we find the asymptotic expansion at O of the unique generalized solution. On the other hand, in the general case when $f \in C^1(\overline{\Omega})$ the problem is not Fredholm because it has an infinite dimensional cokernel. We show that there are an infinite number of necessary conditions for the existence of bounded solutions. We discuss the semi-Fredholm solvability of Problem $P2$ and for $f \in C^6(\overline{\Omega})$ we prove that the necessary conditions for the existence of bounded solutions are also sufficient.

In a historical perspective, Protter studied Problems $P2$ and $P2^*$ in connection with BVPs for mixed type equations that model transonic flow phenomena. In fact, in [33] he also proposes a multidimensional analogue to the two-dimensional Guderley-Morawetz problem for the Gellerstedt equation of hyperbolic-elliptic type. The Guderley-Morawetz problem describes flows around airfoils and is well studied. The existence of a weak solutions and the uniqueness of the strong ones were first established by Morawetz [23] by reducing the problem to a first-order system. Lax and Phillips [20] established that these weak solutions are strong. A survey for the classical 2-D mixed-type BVPs and their transonic background can be found in [24]. The domain of Protter's analogue could be constructed by rotation in \mathbb{R}^4 of a symmetric planar domain for Guderley-Morawetz problem around the axis of symmetry. As a result the set Ω forms the hyperbolic part of the domain. Although it was expected the multidimensional mixed-type problems to be similar to the

two-dimensional BVPs, for the Protter hyperbolic-elliptic problems a general understanding of the situation is still not at hand. Even the question of well posedness is surprisingly subtle and not completely resolved. One has uniqueness results for quasiregular solutions, a class of solutions introduced by Protter, but there are real obstructions to existence in this class. The Protter problems in the hyperbolic part Ω of the domain illustrate some of the difficulties and differences between the planar BVPs and the multidimensional analogues.

In order to construct the solutions of the homogenous Problem $P2^*$ we need the *spherical functions* Y_n^m in \mathbb{R}^3 . Traditionally, Y_n^m are defined on the unit sphere $S^2 := \{(x_1, x_2, x_3) : x_1^2 + x_2^2 + x_3^2 = 1\}$ (see [14]). For convenience in the discussions that follows, we keep the same notation Y_n^m for the radial extension of the spherical function to $\mathbb{R}^3 \setminus \{O\}$, i.e. $Y_n^m(x) := Y_n^m(x/|x|)$ for $x \in \mathbb{R}^3 \setminus \{O\}$. For the definition and properties of the spherical functions see Section 3. For $n, k \in \mathbb{N} \cup \{0\}$ define the functions

$$(1.2) \quad E_k^n(x, t) = \sum_{i=0}^k B_{k,i}^n \frac{(|x|^2 - t^2)^{n-k-i}}{|x|^{n-2i+1}},$$

where the coefficients are

$$B_{k,i}^n := (-1)^i \frac{(k-i+1)_i (n+1-k-i)_i}{i! (n-i+\frac{1}{2})_i}, \quad B_{k,0}^n = 1,$$

with $(a)_i := a(a+1) \cdots (a+i-1)$ and $(a)_0 := 1$. Then the functions

$$W_{k,m}^n(x, t) = E_k^n(x, t) Y_n^m(x)$$

are classical solutions of the homogeneous adjoint Protter problems.

Lemma 1.1. [29] *The functions $W_{k,m}^n(x, t)$ are classical solutions from $C(\overline{\Omega}) \cap C^\infty(\overline{\Omega} \setminus \{O\})$ of the homogeneous Problem $P2^*$ for $k = 0, 1, \dots, [n/2] - 1$.*

A necessary condition for the existence of classical solution for the Problem $P2$ is the orthogonality with respect to the $L^2(\Omega)$ inner product, of the right-hand side function f to all functions $W_{k,m}^n(x, t)$ from Lemma 1.1. To avoid an infinite number of necessary conditions in the framework of classical solvability, we introduce *generalized solutions* for the Problem $P2$ (see the similar definition for the (2+1)-D case in [27] and [29]).

Definition 1.1. *A function $u = u(x, t)$ is called a generalized solution of the Problem $P2$ in Ω , if the following conditions are satisfied:*

- 1) $u \in C^1(\overline{\Omega} \setminus O)$, $u_t|_{\Sigma_0 \setminus O} = 0$, $u|_{\Sigma_1} = 0$, and
- 2) the identity

$$\int_{\Omega} (u_t w_t - u_{x_1} w_{x_1} - u_{x_2} w_{x_2} - u_{x_3} w_{x_3} - f w) dx dt = 0$$

holds for all $w \in C^1(\overline{\Omega})$ such that $w_t|_{\Sigma_0} = 0$ and $w = 0$ in a neighborhood of Σ_2 .

This definition allows the generalized solution to have singularity at the origin and we have an uniqueness result (see Theorem 5.1). Without any additional conditions imposed on the right-hand side function $f \in C(\bar{\Omega})$, it is known (see [26, 29]) that the generalized solution may have power type singularity. Alternatively, we will prove the following necessary conditions for the existence of bounded solutions.

Theorem 1.1. *Suppose that there is a bounded generalized solution of the Protter Problem P2 with right-hand side function $f(x, t) \in C(\bar{\Omega})$. Then*

$$(1.3) \quad \int_{\Omega} W_{k,m}^n(x, t) f(x, t) \, dx dt = 0,$$

for all $n \in \mathbb{N} \cup \{0\}$, $k = 0, \dots, [n/2]$, $m = 1, \dots, 2n + 1$.

The proof of Theorem 1.1 is given in Section 4, but before that we will describe the exact influence of the conditions (1.3) on the behavior of the generalized solution.

First, we consider the case when the right-hand side function $f \in C^1(\bar{\Omega})$ of the wave equation (1.1) has the representation

$$(1.4) \quad f(x, t) = \sum_{n=0}^l \sum_{m=1}^{2n+1} f_n^m(|x|, t) Y_n^m(x),$$

with $l \in \mathbb{N} \cup \{0\}$. In particular, notice that in the case when $f_n^m(|x|, t) = |x|^n a_n^m(t)$ the function f is a harmonic polynomial in x of order l , whose coefficients are functions of t (see the properties of Y_n^m in Section 3). For convenience further by ‘‘harmonic polynomial of order l ’’ we will mean a function from $C^1(\bar{\Omega})$ that has the more general form (1.4). The coefficients f_n^m are

$$f_n^m(|x|, t) = \int_{S^2} f Y_n^m \, d\sigma$$

and must have some special properties at $(0, 0)$ (see for example Lemma 3.3).

According to the results from [29] we know that the *generalized solution* of Problem P2 may have a power type singularity at the origin $O : x = 0, t = 0$. In the present paper we study more accurately the exact behavior of the solution of Problem P2 at O . It is governed by the parameters

$$(1.5) \quad \beta_{k,m}^n := \int_{\Omega} W_{k,m}^n(x, t) f(x, t) \, dx dt,$$

where $n = 0, \dots, l$; $k = 0, \dots, [\frac{n}{2}]$ and $m = 1, \dots, 2n + 1$. We find the asymptotic formula for the *generalized solution* of Problem P2.

Theorem 1.2. *Suppose that the right-hand side function $f \in C^1(\bar{\Omega})$ has the form (1.4). Then the unique generalized solution $u(x, t)$ of Problem P2 belongs*

to $C^2(\overline{\Omega} \setminus O)$ and has the following asymptotic expansion at the singular point O

$$u(x, t) = \sum_{p=1}^{l+1} (|x|^2 + t^2)^{-p/2} F_p(x, t) + F(x, t),$$

where:

(i) the function $F \in C^2(\overline{\Omega} \setminus O)$ and satisfies the a priori estimate

$$|F(x, t)| \leq C \|f\|_{C^1(\Omega)}, \quad (x, t) \in \Omega,$$

with constant C independent of f and $\|f\|_{C^k(\Omega)} = \sum_{|\alpha| \leq k} \max_{\overline{\Omega}} |D^\alpha f(x, t)|$;

(ii) the functions F_p , $p = 1, \dots, l+1$, satisfy the equalities

$$(1.6) \quad F_p(x, t) = \sum_{k=0}^{[(l-p+1)/2]} \sum_{m=1}^{2p+4k-1} \beta_{k,m}^{p+2k-1} F_{k,m}^{p+2k-1}(x, t),$$

with functions $F_{k,m}^n \in C^2(\overline{\Omega} \setminus O)$ bounded and independent of f ;

(iii) if at least one of the constants $\beta_{k,m}^{p+2k-1}$ in (1.6) is different from zero, then for the corresponding function $F_p(x, t)$ there exists a direction $(\alpha, 1) := (\alpha_1, \alpha_2, \alpha_3, 1)$ with $(\alpha, 1)t \in \Sigma_2$ for $0 < t < 1/2$, such that

$$\lim_{t \rightarrow +0} F_p(\alpha t, t) = c_{p,\alpha} = \text{const} \neq 0.$$

After the case of the harmonic polynomials, here we deal with the more general situation when the right-hand side function f is smooth, but it can not be expanded simply as a sum (1.4). Now, Lemma 1.1 shows that the Problem P2 is not Fredholm solvable.

Remark 1.1. Consider the operator

$$T : u_f \longmapsto f \in C^k(\overline{\Omega}),$$

where u_f is the unique classical solution to Protter Problem P2 for the right-hand side function f . According to Lemma 1.1 we have $\dim \text{coker}(T) = \infty$. This means that T is not Fredholm operator for example in $C^k(\overline{\Omega})$. On the other hand, the uniqueness result Theorem 5.1 shows that $\dim \ker(T) = 0$ and T could be a semi-Fredholm operator. A semi-Fredholm operator is a bounded operator that has a finite dimensional kernel or cokernel, and closed range (see, for example [25]). Accordingly we need to find the range of T .

The next result suggests that T is a semi-Fredholm operator.

Theorem 1.3. Let the function $f(x, t)$ belong to $C^6(\overline{\Omega})$. Then the necessary and sufficient conditions for existence of bounded generalized solution $u(x, t)$ of the Protter Problem P2 are

$$(1.7) \quad \int_{\Omega} W_{k,m}^n(x, t) f(x, t) dx dt = 0,$$

for all $n \in \mathbb{N} \cup \{0\}$, $k = 0, \dots, [n/2]$, $m = 1, \dots, 2n + 1$.

Moreover, this generalized solution $u(x, t) \in C^1(\overline{\Omega} \setminus O)$ and satisfies the a priori estimates

$$|u(x, t)| \leq C \left(\|f\|_{C^5(\overline{\Omega})} + \|f_t\|_{C^5(\overline{\Omega})} \right);$$

$$\sum_{i=1}^3 |u_{x_i}(x, t)| + |u_t(x, t)| \leq C(|x|^2 + t^2)^{-1} \|f\|_{C^6(\overline{\Omega})}$$

where the constant C is independent of the function $f(x, t)$.

Obviously, the set of all functions from $C^6(\overline{\Omega})$ that satisfy the orthogonality conditions (1.7) is closed. Therefore, Theorem 1.3 shows that the operator T defined in Remark 1.1 with a domain $D(T) \subset C(\overline{\Omega})$ has a closed range in $C^6(\overline{\Omega})$, and we get the following result.

Corollary 1.1. *The operator T is a semi-Fredholm operator from $D(T) \subset C(\overline{\Omega})$ to $C^6(\overline{\Omega})$.*

We have briefly announced some of the results from this section in [32] with the assumption $f \in C^9(\overline{\Omega})$.

The main results in this work are discussed in Section 2 and the proofs are in Sections 3–7. In more detail the paper is organized as follows: estimates for the spherical functions involved in the representation of the solution are proven in Section 3. In Section 4 the necessary conditions for bounded solution Theorem 1.1 is proved. In Section 5 we consider some two dimensional boundary value problems connected to Problem $P2$ – the Problems $P21$ and $P22$. Exact formulas for the solution of the Problem $P22$ are presented in Lemma 5.1. In Section 6 the proofs of the main Theorems 1.2 and 1.3 are given based on the results from the previous sections and an asymptotic expansion formula for the generalized solution of the 2D Problem $P21$ (Theorem 6.1). The long and technical proof of Theorem 6.1 is postponed to the last Section 7.

2. HISTORICAL REMARKS ON THE MAIN RESULTS

Let us point out several related recent works on Protter problems. Necessary and sufficient conditions for the existence of solutions with fixed order of singularity were obtained in [29]. Similarly, for the \mathbb{R}^3 -analogues of Protter problems some results are presented in [7, 28]. For the problem with Dirichlet type boundary condition on Σ_0 , formula for the asymptotic expansion of the singular solution can be found in [30], and the semi-Fredholm solvability is discussed in [31] for $f \in C^{10}(\overline{\Omega})$. A comparison of various recent results for Protter problems is made in [7].

Various authors adopted a variety of approaches to Protter problems over the last sixty years, for example: Wiener-Hopf method, special Legendre functions, a priori estimates, nonlocal regularization, etc. (see [27] and references there, see also [2, 8, 15, 19, 28, 29]). Alternatively, different multidimensional analogues of the classical Darboux problem for the wave equation are considered in [4, 5, 17], while for some related semilinear equations and systems see [18]. The existence of bounded or unbounded solutions for the wave equation in \mathbb{R}^3 and \mathbb{R}^4 , as well

as for the Euler-Poisson-Darboux equation has been studied in [1, 2, 19, 15, 16, 12, 29].

Regarding the Protter problems with lower order terms see [11] and references therein. Problems with more general boundary condition $u_t + \alpha(x)u = 0$ on Σ_0 are studied in [11, 12]. Some possible regularization methods involving integrodifferential or nonlocal terms can be found in [8].

For the Protter problems for equations of mixed hyperbolic-elliptic type proposed in [33], Aziz and Schneider [3] proved an uniqueness result in the linear case (see also [5]). Concerning nonexistence principle for nontrivial solution of semilinear mixed-type equations in multidimensional case, we refer to [22].

In 1960 Garabedian [10] proved the uniqueness of a classical solution of Protter problem. However, generally, Problem $P2$ is not classically solvable and a necessary condition for the existence of a classical solution is the orthogonality of the right-hand side function f to all solutions of the corresponding homogeneous adjoint Problem $P2^*$. Here in Lemma 1.1, the solutions $W_{k,m}^n$ were constructed with the help of the functions E_k^n defined by (1.2). The alternate representation in terms of the Gauss hypergeometric function $F = F(a, b, c; x)$

$$E_k^n(x, t) = |x|^{n-2k-1}(1 - t^2/|x|^2)^{n-2k}F(n - k + 1/2, -k, 1/2; t^2/|x|^2)$$

can be found in Khe Kan Cher [19]. In [26] there are some solutions for the three-dimensional analogue of the homogeneous Problem $P2^*$.

Let us look back at Theorem 1.1 and the necessary orthogonality conditions (1.3) for the existence of bounded solutions of Problem $P2$. Naturally, these conditions include the functions $W_{k,m}^n$ from Lemma 1.1. However, notice that there are also some others.

Remark 2.1. *It is interesting that the conditions (1.3) include the case of even $n = 2k$. Notice that the functions $W_{k,m}^{2k}(x, t)$ are not classical solutions of the homogenous adjoint Problem $P2^*$. Actually, they satisfy the homogenous wave equation in Ω and $(W_{k,m}^{2k})_t$ vanish on Σ_0 , but $W_{k,m}^{2k}$ is not zero on Σ_2 . In addition, the functions $W_{k,m}^{2k}$ have a singularity at the origin O like $|x|^{-1}$, however this singularity is integrable in the domain Ω .*

Instead of imposing an infinite number of orthogonality conditions on f , Popivanov and Schneider [26, 27] introduced the concept of generalized solution that allows the solution to have singularity on the inner cone Σ_2 . Here, Theorem 1.2 describes the affect of the parameters $\beta_{k,m}^n$ on the behavior of the *generalized solution* of Problem $P2$. The constants $\beta_{k,m}^n$ are defined by (1.5) and are obviously related to the orthogonality conditions (1.3). When the right-hand function is a harmonic polynomial (1.4), the asymptotic expansion in Theorem 1.2 shows that the *generalized solution* could be bounded only if all $\beta_{k,m}^n$ involved are zero.

Corollary 2.1. *Suppose that the right-hand side function $f \in C^1(\overline{\Omega})$ has the form (1.4) and satisfies the orthogonality conditions*

$$(2.1) \quad \int_{\Omega} W_{k,m}^n(x,t) f(x,t) \, dx dt = 0$$

for all $n = 0, \dots, l$; $k = 0, \dots, \lfloor \frac{n}{2} \rfloor$ and $m = 1, \dots, 2n + 1$. Then the unique generalized solution $u(x,t)$ of Problem P2 belongs to $C^2(\overline{\Omega} \setminus O)$, is bounded and satisfies the a priori estimate

$$\max_{\overline{\Omega}} |u| \leq C \|f\|_{C^1(\Omega)}.$$

On the other hand, without any orthogonality conditions on f , the following result is obtained.

Corollary 2.2. *The generalized solution u of Problem P2 with a right-hand side function $f \in C^1(\overline{\Omega})$ in the form (1.4) satisfies the a priori estimate*

$$(2.2) \quad |u(x,t)| \leq C \left(\max_{\overline{\Omega}} |f| \right) (|x|^2 + t^2)^{-(l+1)/2}.$$

The influence of the orthogonality conditions (2.1) on the exact behavior of the *generalized solution* is clarified by the case (iii) of Theorem 1.2. It shows that for fixed indexes (n, k, m) , the corresponding condition (2.1) ‘‘controls’’ one power-type singularity.

In the Corollaries 2.1 and 2.2 the emphasis is on the extreme cases: when all orthogonality conditions (2.1) are fulfilled or, alternatively, when none of them are satisfied. In both cases the exact behavior of the solution is given. The estimate (2.2), presented here is analogous to known estimates for Protter problems in \mathbb{R}^3 ([27]) and in \mathbb{R}^m ([1]). It is interesting that singularities of the *generalized solutions* are isolated at the origin and do not propagate in the direction of the bicharacteristics on the characteristic cone Σ_2 . Traditionally, it is assumed that the wave equation, with sufficiently smooth right-hand side cannot have a solution with an isolated singular point as in Hörmander [13, Chapter 24.5]. The case here is different since the point of singularity O lies on the non-characteristic part of the boundary Σ_0 as well as on the characteristic part Σ_2 .

Remark 2.2. *The Problem P2 in \mathbb{R}^4 with harmonic polynomial on the right-hand side is also studied in [29]. However, the explicit asymptotic expansion here has no analogue in [29], where only the behavior of the singularities is given. Additionally, if the orthogonality conditions (2.1) are fulfilled, the Corollary 2.1 states that the generalized solution is in fact bounded, while the estimates in [29, Theorem 1.1] still allow the solution to have some logarithmic singularities.*

Remark 2.3. *Let us compare Protter problems in \mathbb{R}^3 (as treated in [28, 7]) and \mathbb{R}^4 here (see also [29, 30]). In both cases the study these BVPs is based on the properties of the special Legendre functions. Instead of Legendre polynomials P_n here, in the three-dimensional case the Legendre functions P_ν with non-integer*

indexes $\nu = n - 1/2$ are used (for their properties see [9]). One can easily modify both these techniques to obtain similar results for the $(m+1)$ -dimensional problems: for even m (analogous to \mathbb{R}^3 case) or for odd m (the present case \mathbb{R}^4). Some related results for Protter problems in \mathbb{R}^{m+1} are presented in [1, 2].

In the general case when the right-hand side function f is smooth enough, Theorem 1.3 implies that the necessary conditions (1.3) for existence of bounded solutions from Theorem 1.1 are also sufficient. Further, this means that there are no other nontrivial classical solutions of the homogenous adjoint Problem $P2^*$ except those listed in Lemma 1.1.

Remark 2.4. We point out the differences between Theorem 1.3 and the results from [31] for Protter Problem P1 with Dirichlet type boundary condition on Σ_0 . First, notice that in the case when right-hand side function f is harmonic polynomial of order l the solution of Problem P2 may have worse singularity (like $(|x|^2 + t^2)^{-(l+1)/2}$, see Theorem 1.2) than the solution of Problem P1 (like $(|x|^2 + t^2)^{-l/2}$, see [30]). For the general case we are able to reduce the assumptions on f – in Theorem 1.3 we assume $f \in C^6(\bar{\Omega})$, while in [31, Theorem 1.1] smoother $f \in C^{10}$ is required. In order to achieve this, we rely on the more accurate estimates for the special functions proven in Section 3.

3. ESTIMATES FOR THE SPECIAL FUNCTIONS

For the proof of the main results we will need some properties of the *spherical functions* Y_n^m in \mathbb{R}^3 . They are naturally expressed on the unit sphere $S^2 := \{(x_1, x_2, x_3) : x_1^2 + x_2^2 + x_3^2 = 1\}$ in spherical polar coordinates. Let us introduce polar coordinates (r, θ, φ) in \mathbb{R}^3 :

$$x_1 = r \sin \theta \cos \varphi, \quad x_2 = r \sin \theta \sin \varphi, \quad x_3 = r \cos \theta.$$

where $0 \leq \theta < \pi$, $0 \leq \varphi < 2\pi$, $r > 0$. Then the spherical functions, expressed in terms of θ and φ as in the traditional definition on S^2 (see [14]), are given by

$$Y_n^{2k+1}(\theta, \varphi) = \sqrt{\frac{2n+1}{2\pi}} \sqrt{\frac{(n-k)!}{(n+k)!}} P_n^k(\cos \theta) \cos k\varphi, \quad k = 1, \dots, n$$

(3.1)

$$Y_n^{2k}(\theta, \varphi) = \sqrt{\frac{2n+1}{2\pi}} \sqrt{\frac{(n-k)!}{(n+k)!}} P_n^k(\cos \theta) \sin k\varphi, \quad k = 1, \dots, n,$$

and $Y_n^1(\theta, \varphi) = ((2n+1)/4\pi)^{1/2} P_n(\cos \theta)$. Here P_n are the *Legendre polynomials* defined by the Rodrigues' formula

$$P_n(s) := \frac{1}{2^n n!} \frac{d^n}{ds^n} (s^2 - 1)^n = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} s^{n-2k}, \quad a_{2k} \neq 0.$$

(3.2)

while P_n^k are the *associated Legendre polynomials* that can be defined as

$$P_n^k(s) = (-1)^k (1-s^2)^{k/2} \frac{d^k}{ds^k} P_n(s).$$

The functions Y_n^m satisfy the differential equation

$$(3.3) \quad \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y_n^m \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} Y_n^m + n(n+1)Y_n^m = 0.$$

and form a complete orthonormal system in $L_2(S^2)$ (see [14]).

Using Cartesian coordinates as in Section 1, one can define the spherical functions as $Y_n^m(x_1, x_2, x_3) := Y_n^m(\theta, \varphi)$ for $x \in S^2$, or by

$$Y_n^{2m}(x_1, x_2, x_3) = C_n^m \frac{d^m}{dx_3^m} P_n(x_3) \operatorname{Im} \{(x_2 + ix_1)^m\}, \text{ for } m = 1, \dots, n$$

and

$$Y_n^{2m+1}(x_1, x_2, x_3) = C_n^m \frac{d^m}{dx_3^m} P_n(x_3) \operatorname{Re} \{(x_2 + ix_1)^m\}, \text{ for } m = 0, \dots, n,$$

where C_n^m are constants. In the present paper, we keep the same notation Y_n^m for the radial extension of the spherical function to $\mathbb{R}^3 \setminus \{O\}$, i.e. $Y_n^m(x) := Y_n^m(x/|x|)$ for $x \in \mathbb{R}^3 \setminus \{O\}$. According to the properties of $Y_n^m(x)$, the function $|x|^n Y_n^m(x)$ is a homogenous harmonic polynomial of order n in the variables x_1, x_2, x_3 .

We will need some estimates for Y_n^m and the special functions involved in the representations of the solutions of the Protter problems. Let us start with the Legendre polynomials P_n .

Lemma 3.1. *The following estimates hold for $x \in [0, 1]$:*

$$(3.4) \quad |P_n(x)| \leq 1, \quad |P_n'(x)| \leq \frac{n(n+1)}{2},$$

$$(3.5) \quad |P_n(x) - P_n(0)| \leq nx.$$

Proof. The estimates (3.4) are proved for example in [29]. Here we will show that (3.5) holds. Using Bonnet's recursion formula

$$(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$$

we get $(n+1)P_{n+1}(0) = -nP_{n-1}(0)$ and

$$(n+1)(P_{n+1}(x) - P_{n+1}(0)) = (2n+1)xP_n(x) - n(P_{n-1}(x) - P_{n-1}(0)).$$

Thus

$$|P_{n+1}(x) - P_{n+1}(0)| \leq 2x + |P_{n-1}(x) - P_{n-1}(0)|.$$

From here and the equalities $P_0(x) - P_0(0) = 0$ and $P_1(x) - P_1(0) = x$ we get the estimate (3.5) by induction. \square

In order to study the first derivatives of the generalized solution of the Protter problem, we will need to estimate also the first derivatives with respect to x of $Y_n^m(x) := Y_n^m(x/|x|)$ – the radial extension of the spherical function to $\mathbb{R}^3 \setminus \{O\}$:

$$(3.6) \quad \begin{aligned} \frac{\partial}{\partial x_1} Y_n^m &:= r^{-1} [\cos \theta \cos \varphi (Y_n^m)_\theta - (\sin \theta)^{-1} \sin \varphi (Y_n^m)_\varphi], \\ \frac{\partial}{\partial x_2} Y_n^m &= r^{-1} [\cos \theta \sin \varphi (Y_n^m)_\theta + (\sin \theta)^{-1} \cos \varphi (Y_n^m)_\varphi], \end{aligned}$$

$$\frac{\partial}{\partial x_3} Y_n^m := -r^{-1} \sin \theta (Y_n^m)_\theta.$$

Using the so called addition theorem for Legendre polynomials we get the following result.

Lemma 3.2. *For $n \geq 1$ the functions $Y_n^m(x)$ satisfy the equalities*

$$\sum_{m=1}^{2n+1} (Y_n^m)^2 = \frac{2n+1}{4\pi},$$

$$\sum_{m=1}^{2n+1} \sum_{i=1}^3 \left(\frac{\partial}{\partial x_i} Y_n^m \right)^2 = \frac{n(n+1)(2n+1)}{2\pi|x|^2}.$$

Proof. From the definition (3.1) of Y_n^m follows that

$$\sum_{m=1}^{2n+1} (Y_n^m)^2 = \frac{2n+1}{4\pi} \left[(P_n(\cos \theta))^2 + 2 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} (P_n^k(\cos \theta))^2 \right]$$

According to the Addition theorem (see [21])

$$(3.7) \quad \begin{aligned} & P_n(\cos \theta \cos \theta_1 + \sin \theta \sin \theta_1 \cos \varphi) \\ &= P_n(\cos \theta) P_n(\cos \theta_1) + 2 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} P_n^k(\cos \theta) P_n^k(\cos \theta_1) \cos k\varphi. \end{aligned}$$

With $\theta = \theta_1$ and $\varphi = 0$ one derives the equality

$$(P_n(\cos \theta))^2 + 2 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} (P_n^k(\cos \theta))^2 = 1.$$

This means that

$$\sum_{m=1}^{2n+1} (Y_n^m)^2 = \frac{2n+1}{4\pi}.$$

Using (3.7) again, we get the required property of the derivatives of Y_n^m . Directly from (3.6) for the squares of the derivatives with respect to x we have

$$(3.8) \quad \sum_{|\alpha|=1} (D_x^\alpha Y_n^m)^2 = r^{-2} (Y_n^m)_\theta^2 + r^{-2} (\sin \theta)^{-2} (Y_n^m)_\varphi^2$$

and from the definition (3.1) of Y_n^m we find

$$\sum_{m=1}^{2n+1} \left(\frac{\partial Y_n^m}{\partial \theta} \right)^2 = \frac{2n+1}{4\pi} \left[\left(\frac{\partial}{\partial \theta} P_n(\cos \theta) \right)^2 + 2 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} \left(\frac{\partial}{\partial \theta} P_n^k(\cos \theta) \right)^2 \right];$$

$$\sum_{m=1}^{2n+1} \left(\frac{\partial Y_n^m}{\partial \varphi} \right)^2 = \frac{2n+1}{2\pi} \sum_{k=1}^n k^2 \frac{(n-k)!}{(n+k)!} (P_n^k(\cos \theta))^2.$$

Put $\varphi = 0$ and differentiation of (3.7) with respect to θ and θ_1 gives

$$\begin{aligned} & P_n''(\cos \theta \cos \theta_1 + \sin \theta \sin \theta_1) \{ \cos \theta \sin \theta_1 - \sin \theta \cos \theta_1 \} \{ \sin \theta \cos \theta_1 - \cos \theta \sin \theta_1 \} \\ & \quad + P_n'(\cos \theta \cos \theta_1 + \sin \theta \sin \theta_1) \{ \cos \theta \cos \theta_1 + \sin \theta \sin \theta_1 \} \\ & = \sin \theta \sin \theta_1 P_n'(\cos \theta) P_n'(\cos \theta_1) + 2 \sin \theta \sin \theta_1 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} P_n^{k'}(\cos \theta) P_n^{k'}(\cos \theta_1). \end{aligned}$$

With $\theta = \theta_1$ we derive

$$\left(\frac{\partial}{\partial \theta} P_n(\cos \theta) \right)^2 + 2 \sum_{k=1}^n \frac{(n-k)!}{(n+k)!} \left(\frac{\partial}{\partial \theta} P_n^k(\cos \theta) \right)^2 = P_n'(1).$$

Analogously, after differentiating (3.7) twice with respect to φ :

$$\begin{aligned} & P_n''(\cos \theta \cos \theta_1 + \sin \theta \sin \theta_1 \cos \varphi) \sin^2 \theta \sin^2 \theta_1 \sin^2 \varphi \\ & \quad - P_n'(\cos \theta \cos \theta_1 + \sin \theta \sin \theta_1 \cos \varphi) \sin \theta \sin \theta_1 \cos \varphi \\ & = -2 \sum_{k=1}^n k^2 \frac{(n-k)!}{(n+k)!} P_n^k(\cos \theta) P_n^k(\cos \theta_1) \cos k\varphi. \end{aligned}$$

Then substituting $\varphi = 0$ and $\theta_1 = \theta$ we have

$$2 \sum_{k=1}^n k^2 \frac{(n-k)!}{(n+k)!} (P_n^k(\cos \theta))^2 = (\sin \theta)^2 P_n'(1).$$

Since $P_n'(1) = n(n+1)/2$ we conclude that

$$\sum_{m=1}^{2n+1} \sum_{|\alpha|=1} (D_x^\alpha Y_n^m)^2 = \frac{n(n+1)(2n+1)}{2\pi r^2}.$$

□

As a direct consequence, we find the next estimates for the spherical functions radially extended out of S^2 .

Corollary 3.1. *For $n \geq 1$ the functions $Y_n^m(x)$ satisfy the inequalities*

$$(3.9) \quad |Y_n^m(x)| \leq n^{1/2},$$

$$(3.10) \quad \sum_{m=1}^{2n+1} |Y_n^m(x)| \leq n,$$

$$(3.11) \quad \sum_{m=1}^{2n+1} \sum_{i=1}^3 \left| \frac{\partial}{\partial x_i} Y_n^m(x) \right| \leq 3n^2 |x|^{-1},$$

$$(3.12) \quad \sum_{m=1}^{2n+1} [|(Y_n^m)_\theta| + |(\sin \theta)^{-1} (Y_n^m)_\varphi|] \leq 3n^2.$$

Proof. Obviously

$$|Y_n^m(x)| \leq \left[\sum_{m=1}^{2n+1} (Y_n^m)^2 \right]^{1/2} \leq n^{1/2}.$$

The estimates (3.10) and (3.11) follow directly from the Cauchy-Schwarz inequality and Lemma 3.2

$$\sum_{m=1}^{2n+1} |Y_n^m(x)| \leq \sqrt{2n+1} \left[\sum_{m=1}^{2n+1} (Y_n^m)^2 \right]^{1/2} = \frac{2n+1}{2\sqrt{\pi}} \leq n,$$

$$\sum_{m=1}^{2n+1} \sum_{i=1}^3 \left| \frac{\partial}{\partial x_i} Y_n^m(x) \right| \leq \sqrt{6n+3} \left[\sum_{m=1}^{2n+1} \sum_{i=1}^3 \left(\frac{\partial}{\partial x_i} Y_n^m \right)^2 \right]^{1/2} = \frac{(2n+1)\sqrt{3n(n+1)}}{\sqrt{2\pi}|x|}.$$

Finally (3.12) follows from the equality (3.8):

$$\begin{aligned} \sum_{m=1}^{2n+1} \{ |(Y_n^m)_\theta| + |(\sin \theta)^{-1}(Y_n^m)_\varphi| \} &\leq \sqrt{4n+2} \left\{ \sum_{m=1}^{2n+1} [(Y_n^m)_\theta]^2 + (\sin \theta)^{-2} (Y_n^m)_\varphi^2 \right\}^{1/2} \\ &= |x| \sqrt{4n+2} \left[\sum_{m=1}^{2n+1} \sum_{i=1}^3 \left(\frac{\partial}{\partial x_i} Y_n^m \right)^2 \right]^{1/2} \leq 3n^2. \end{aligned}$$

□

Generally, if the function $f(x, t)$ is smooth enough, it can be represented as a harmonic series. In order to estimate the coefficients of the series we will use the next result. It is based on the fact that the spherical functions are eigenfunctions for the Laplace operator on the sphere S^2 .

Lemma 3.3. *Let $k \in \mathbb{N} \cup \{0\}$.*

1) ([31]) *Let $f \in C^{2k}(\bar{\Omega})$. Then it has the representation*

$$f(x, t) = \sum_{n=0}^{\infty} \sum_{m=1}^{2n+1} f_n^m(|x|, t) Y_n^m(x)$$

and for $n \in \mathbb{N}$

$$|f_n^m| \leq C n^{-2k} \|f\|_{C^{2k}(\bar{\Omega})}.$$

2) *Suppose $f \in C^{2k+1}(\bar{\Omega})$ then for $n \in \mathbb{N}$*

$$\sum_{m=1}^{2n+1} |f_n^m| \leq C n^{-2k} |x| \|f\|_{C^{2k+1}(\bar{\Omega})}.$$

Proof. The estimate in the case 1) follows directly from the fact that the spherical functions are eigenfunctions for the spherical Laplacean Δ_S

$$\Delta_S F := \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} F \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} F,$$

i.e., the equation (3.3) shows that $\Delta_S Y_n^m = -n(n+1)Y_n^m$. In fact f_n^m is bounded by the Fourier coefficient of $(\Delta_S)^k f$:

$$|f_n^m| = \left| \int_{S^2} Y_n^m f \, d\sigma \right| \leq C n^{-2k} \left| \int_{S^2} Y_n^m (\Delta_S)^k f \, d\sigma \right|.$$

For detail proof of 1) see [31] and [35].

Next we will prove the case 2) with $k = 0$. Let $F \in C^1(\bar{\Omega})$ then

$$F_n^m(r, t) := \int_{S^2} Y_n^m F \, d\sigma = -n^{-1}(n+1)^{-1} \int_0^\pi \left(\int_0^{2\pi} (\Delta_S Y_n^m) F \, d\varphi \right) \sin \theta \, d\theta.$$

Integrating by parts gives

$$F_n^m = \frac{1}{n(n+1)} \int_0^\pi \left[\int_0^{2\pi} \left(\sin \theta \frac{\partial}{\partial \theta} Y_n^m \frac{\partial}{\partial \theta} F + \frac{1}{\sin \theta} \frac{\partial}{\partial \varphi} Y_n^m \frac{\partial}{\partial \varphi} F \right) d\varphi \right] d\theta,$$

where

$$\begin{aligned} \sin \theta \frac{\partial}{\partial \theta} F &= r \sin \theta \cos \theta \cos \varphi \frac{\partial}{\partial x_1} F + r \sin \theta \cos \theta \sin \varphi \frac{\partial}{\partial x_2} F - r \sin^2 \theta \frac{\partial}{\partial x_3} F \\ \frac{\partial}{\partial \varphi} F &= -r \sin \theta \sin \varphi \frac{\partial}{\partial x_1} F + r \sin \theta \cos \varphi \frac{\partial}{\partial x_2} F. \end{aligned}$$

Therefore using (3.12) we have

$$\begin{aligned} \sum_{m=1}^{2n+1} |F_n^m| &\leq \frac{C|x|}{n^2} \|F\|_{C^1(\bar{\Omega})} \sum_{m=1}^{2n+1} \int_0^\pi \left[\int_0^{2\pi} (|(Y_n^m)_\theta| + |(\sin \theta)^{-1} (Y_n^m)_\varphi|) d\varphi \right] d\theta \\ &\leq C_1 |x| \|F\|_{C^1(\bar{\Omega})}. \end{aligned}$$

To prove the case 2) it remains to substitute $(\Delta_S)^k f \in C^1(\bar{\Omega})$ for F in this estimate. \square

4. NECESSARY CONDITIONS FOR BOUNDED SOLUTION

Here we will prove of the necessity of the orthogonality conditions (1.3) for the existence of a bounded solution.

Proof of Theorem 1.1. Let u be a bounded generalized solution of Problem P2. Let us fix a function $\chi(s) \in C^\infty(\mathbb{R})$ such that $\chi(s) = 0$ for $s \leq 1$, and $\chi(s) = 1$ for $s \geq 2$.

For fixed indexes $n \in \mathbb{N} \cup \{0\}$, $0 \leq k \leq [\frac{n}{2}]$, $1 \leq m \leq 2n+1$, consider the functions

$$W_q(x, t) = \chi(2q(|x| - t)) \chi(q|x|) W_{k,m}^n(x, t)$$

for $q \in \mathbb{N}$. Obviously, $W_q \in C^\infty(\overline{\Omega})$ vanishes on a neighborhood of Σ_2 and $(W_q)_t = 0$ on Σ_0 . Therefore according to Definition 1.1 we have

$$\int_{\Omega} \{u_t(W_q)_t - u_{x_1}(W_q)_{x_1} - u_{x_2}(W_q)_{x_2} - u_{x_3}(W_q)_{x_3}\} dxdt = \int_{\Omega} fW_q dxdt$$

and thus

$$(4.1) \quad \int_{\Omega} fW_q dxdt = \int_{\Omega} u \{(W_q)_{x_1x_1} + (W_q)_{x_2x_2} + (W_q)_{x_3x_3} - (W_q)_{tt}\} dxdt.$$

We want to prove that $\int_{\Omega} fW_{k,m}^n dxdt = 0$ for all $n \in \mathbb{N} \cup \{0\}$, $k = 0, \dots, [n/2]$, $m = 1, \dots, 2n + 1$. Notice that $|fW_{k,m}^n| \leq C|x|^{-1}$ and then the integral $\int_{\Omega} |fW_{k,m}^n| dxdt$ is convergent. The function $f \in C(\overline{\Omega})$ and thus

$$\lim_{q \rightarrow +\infty} \int_{\Omega} fW_q dxdt = \int_{\Omega} fW_{k,m}^n dxdt.$$

Therefore it is sufficient to prove that when $q \rightarrow +\infty$ the right hand side of the equality (4.1) tends to zero for $n \in \mathbb{N} \cup \{0\}$, $k = 0, \dots, [n/2]$, $m = 1, \dots, 2n + 1$.

Using the fact that the functions $W_{k,m}^n$ are solutions of the homogenous wave equation in Ω , straight forward computations show that

$$\begin{aligned} \sum_{i=1}^3 (W_q)_{x_i x_i} - (W_q)_{tt} &= \left[(W_{k,m}^n)_r + (W_{k,m}^n)_t + \frac{1}{r} W_{k,m}^n \right] 4q\chi'_1\chi_2 + 4q^2 W_{k,m}^n \chi'_1\chi'_2 \\ &\quad + \left[(W_{k,m}^n)_r + \frac{1}{r} W_{k,m}^n \right] 2q\chi_1\chi'_2 + q^2 W_{k,m}^n \chi_1\chi''_2 \end{aligned}$$

where $r = |x|$, while χ_1 and χ_2 stand for $\chi(2q(r-t))$ and $\chi(qr)$ respectively. Expressed with spherical polar coordinates (r, φ, θ) in \mathbb{R}^3

$$\begin{aligned} &\int_{\Omega} u [(W_q)_{x_1x_1} + (W_q)_{x_2x_2} + (W_q)_{x_3x_3} - (W_q)_{tt}] dxdt \\ &= \int_{\Omega} \left\{ \left[(E_k^n)_r + (E_k^n)_t + \frac{1}{r} E_k^n \right] 4q\chi'_1\chi_2 + q^2 E_k^n (4\chi'_1\chi'_2 + \chi_1\chi''_2) \right. \\ &\quad \left. + \left[(E_k^n)_r + \frac{1}{r} E_k^n \right] 2q\chi_1\chi'_2 \right\} u Y_n^m(\theta, \varphi) r^2 \sin \theta dr d\varphi d\theta dt. \end{aligned}$$

For simplicity, we consider only one of the terms from the definition of the function $E_k^n(r, t)$ (see (1.2)). Let us denote by

$$w(r, t) = \frac{(r^2 - t^2)^{n-k-i}}{r^{n-2i+1}}$$

defined in $D = \{0 < t < 1/2, t < r < 1 - t\}$, and consider the subsets $D_{1,q} = D \cap \{1/(2q) < r-t < 1/q\} \cap \{r > 1/q\}$ and $D_{2,q} = D \cap \{1/q < r < 2/q\}$.

Since the functions u , χ' , χ'' and Y_n^m are bounded, the function $\chi'(qr) = 0$ in $D \setminus D_{2,q}$ and $\chi'(2q(r-t))\chi(qr)$ is zero in $D \setminus D_{1,q}$, it is sufficient to prove that

the integrals

$$I_{1,q} := q \int_{D_{1,q}} |w_r + w_t + r^{-1}w| r^2 dr dt ;$$

$$I_{2,q} := q \int_{D_{2,q}} |w_r + r^{-1}w| r^2 dr dt$$

and

$$I_{3,q} := q^2 \int_{D_{2,q}} w r^2 dr dt$$

tend to zero as $q \rightarrow \infty$, for $n \in \mathbb{N} \cup \{0\}$, $k = 0, \dots, [n/2]$ and $i = 0, \dots, k$.

We get $\lim_{q \rightarrow \infty} I_{3,q} = 0$ from the estimate $|w| \leq r^{n-2k-1}$ in D and thus

$$I_{3,q} \leq q^2 \int_{r=0}^{r=2/q} \int_{t=0}^{t=r} r^{n-2k+1} dt dr \leq q^2 C \int_0^{2/q} r^2 dr \leq C_1 q^{-1}.$$

For $I_{2,q}$ let us compute first

$$|w_r + r^{-1}w| r^2 = \frac{(r^2 - t^2)^{n-k-i-1}}{r^{n-2i}} [(n-2k)r^2 + (n-2i)t^2] \leq C,$$

because $n-k-i-1$ could be negative only when $n-2k = n-2i = 0$. Therefore, we have

$$I_{2,q} \leq q \int_{r=0}^{r=2/q} \int_{t=0}^{t=r} C dt dr \leq C_1 q^{-1}$$

and thus $\lim_{q \rightarrow \infty} I_{2,q} = 0$.

Finally, to evaluate $I_{1,q}$ we use the estimate

$$|w_r + w_t + r^{-1}w| r^2 = \frac{(r^2 - t^2)^{n-k-i-1}}{r^{n-2i}} (r-t) |(n-2k)r - (n-2i)t| \leq C \frac{r-t}{r}$$

and find

$$I_{1,q} \leq qC \int_{r=1/q}^{r=1} \int_{t=r-1/q}^{t=r} \frac{r-t}{r} dt dr \leq q^{-1} C_1 \int_{r=1/q}^{r=1} r^{-1} dt dr \leq C_2 q^{-1} \ln q$$

that shows that $\lim_{q \rightarrow \infty} I_{1,q} = 0$. □

5. PREVIOUS RESULTS

In this section we quote some results from [29] that will be essentially the starting point for the proofs of Theorems 1.2 and 1.3. We start with the following uniqueness result.

Theorem 5.1. [29] *The Problem P2 has at most one generalized solution.*

In [29] the right-hand side function f of the wave equation (1.1) is fixed as a harmonic polynomial (1.4). Then the following existence result for the generalized solution is valid.

Theorem 5.2. [29] *Suppose that the right-hand side $f \in C^1(\overline{\Omega})$ has the form (1.4) where $l \in \mathbb{N} \cup \{0\}$. Then, the unique generalized solution $u(x, t)$ of the Problem P2 in Ω exists and has the form*

$$(5.1) \quad u(x, t) = \sum_{n=0}^l \sum_{m=1}^{2n+1} u_n^m(|x|, t) Y_n^m(x) \in C^2(\overline{\Omega} \setminus O).$$

In fact, the function $u_n^m(|x|, t)$ from (5.1) is the solution of a two-dimensional boundary value problem that involves only the corresponding coefficient $f_n^m(|x|, t)$ from (1.4). In order to formulate this BVP, it is natural to introduce polar coordinates (r, θ, φ) in \mathbb{R}^3 : r, θ and φ are such that $0 \leq \theta < \pi, 0 \leq \varphi < 2\pi, r > 0$ and

$$x_1 = r \sin \theta \cos \varphi, \quad x_2 = r \sin \theta \sin \varphi, \quad x_3 = r \cos \theta.$$

In the special case when f has the form

$$f(r, \theta, \varphi, t) = f_n^m(r, t) Y_n^m(\theta, \varphi),$$

according to Theorem 5.2 we may look for a solution of the same form

$$u(r, \theta, \varphi, t) = u_n^m(r, t) Y_n^m(\theta, \varphi).$$

Then we can reduce the (3+1)- D Protter problem to some BVPs in \mathbb{R}^2 . From the properties of the spherical functions it follows that the function $u_n^m(r, t)$ is a solution of the equation

$$(5.2) \quad u_{rr} + \frac{2}{r} u_r - u_{tt} - \frac{n(n+1)}{r^2} u = f(r, t)$$

with right-hand side $f(r, t) := f_n^m(r, t)$, in the domain $D = \{(r, t) : 0 < t < 1/2, t < r < 1 - t\}$, bounded by

$$S_0 = \{(r, t) : t = 0, 0 < r < 1\},$$

$$S_1 = \{(r, t) : 0 < t < 1/2, r = 1 - t\}, \quad S_2 = \{(r, t) : 0 < t < 1/2, r = t\}.$$

Thus, we arrive at the next two dimensional problems.

Problem P21. Find a solution of the equation (5.2) in the domain D which satisfies the boundary conditions

$$P21 : \quad u_t|_{S_0} = 0, \quad u|_{S_1} = 0.$$

Finally, we substitute

$$v = ru(r, t), \quad g = rf(r, t)$$

and

$$\xi = \frac{r+t}{2}, \quad \eta = \frac{r-t}{2}$$

and get the following problem.

Problem P22. Find a solution of the equation

$$(5.3) \quad v_{\xi\eta} - \frac{n(n+1)}{(\xi+\eta)^2} v = g$$

in the domain $D_1 = \{(\xi, \eta) \in \mathbb{R}^2 : 0 < \eta < \xi < 1/2\}$ with boundary conditions

$$P22 : \quad \left(\frac{\partial v}{\partial \xi} - \frac{\partial v}{\partial \eta} \right) (\eta, \eta) = 0, \quad v(1/2, \eta) = 0, \quad \text{for } \eta \in (0, 1/2].$$

In [29] the solution of Problem *P22* is constructed with the help of the Riemann's function

$$R(\xi_1, \eta_1; \xi, \eta) = P_n \left(\frac{(\xi - \eta)(\xi_1 - \eta_1) + 2\xi_1\eta_1 + 2\xi\eta}{(\xi_1 + \eta_1)(\xi + \eta)} \right)$$

for the equation (5.3) found by Copson [6]. The problem is reduced to an integral equation of Volterra type. Then this integral equation is solved using some formulas from the book by Samko, Kilbas and Marichev [34] and the properties of the Melin transform. According to formulas (23), (24) and (27) in the proof of Theorem 3.1 from [29] we can write down the following result.

Lemma 5.1. *The solution $v(\xi, \eta)$ of Problem *P22* is given by*

$$(5.4) \quad v(\xi, \eta) = \tau(\xi) + \int_{\xi}^{\frac{1}{2}} \tau(\xi_1) \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 \\ - \int_{\xi}^{\frac{1}{2}} \left(\int_0^{\eta} P_n \left(\frac{(\xi - \eta)(\xi_1 - \eta_1) + 2\xi_1\eta_1 + 2\xi\eta}{(\xi_1 + \eta_1)(\xi + \eta)} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1,$$

where

$$(5.5) \quad \tau(\xi) = \int_{\frac{1}{2}}^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) G(\xi_1) d\xi_1.$$

and

$$(5.6) \quad G(\xi) := \int_{\xi}^{\frac{1}{2}} \left(\int_0^{\xi} P_n \left(\frac{\xi_1\eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1 \\ - \int_0^{\xi} P_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) g \left(\frac{1}{2}, \eta_1 \right) d\eta_1 - \int_{\xi}^{\frac{1}{2}} P_n \left(\frac{\xi}{\xi_1} \right) g(\xi_1, 0) d\xi_1.$$

Finally, we will need the relation between the functions $E_i^n(r, t) := E_i^n(x, t)$, with $r = |x|$, defined in (1.2) and the Legendre polynomials P_n .

Lemma 5.2. [29] *Define the functions*

$$h_k(\xi, \eta) = \int_{\eta}^{\xi} s^k P_n \left(\frac{\xi\eta + s^2}{s(\xi + \eta)} \right) ds.$$

Then the equality

$$r^{-1} \frac{\partial}{\partial t} \left[h_{n-2i} \left(\frac{r+t}{2}, \frac{r-t}{2} \right) \right] = c_i^n E_i^n(r, t)$$

holds for $i = 0, \dots, \lfloor \frac{n}{2} \rfloor$ with some nonzero constants c_i^n .

6. PROOFS OF THE MAIN RESULTS

In this section we will prove Theorem 1.2 and Theorem 1.3. The proofs are based on an asymptotic expansion formula for the solution of the two-dimensional Problem $P21$. In order to formulate it, let us concentrate first on Problem $P2$ with right-hand side functions f of the form

$$f(x, t) = f_n^m(r, t) Y_n^m(\theta, \varphi),$$

with fixed $n, m \in \mathbb{N} \cup \{0\}$ and $m \leq 2n + 1$. We will use the results stated in Section 5. The unique *generalized solution* u of Problem $P2$ also has the form

$$u(x, t) = u_n^m(r, t) Y_n^m(\theta, \varphi).$$

The function $u_n^m(r, t)$ is the solution of Problem $P21$ with the function $f(r, t) := f_n^m(r, t)$ as a right-hand side in the equation (5.2). We are interested in the exact behavior of $u_n^m(r, t)$ at $(0, 0)$. One expects it to depend on the constants

$$(6.1) \quad \beta_k^n := \int_0^{\frac{1}{2}} \left(\int_t^{1-t} E_k^n(r, t) f(r, t) r^2 dr \right) dt \quad \text{for } k = 0, \dots, \lfloor \frac{n}{2} \rfloor,$$

that correspond to $\beta_{k,m}^n$ defined in (1.5) for the Problem $P2$. For simplicity, denote further

$$\begin{aligned} A_0 &:= \|f\|_{C(D)}; \quad A_1 := \|f\|_{C(D)} + \|rf_t\|_{C(D)}; \quad A_2 := \|rf\|_{C^1(D)} + \|rf_t\|_{C^1(D)}; \\ A_3 &:= \|rf\|_{C^1(D)} + \|rf_t\|_{C^1(D)} + \|rf_{tt}\|_{C^1(D)} + \|r^2 f_{trr}\|_{C(D)}. \end{aligned}$$

Theorem 6.1. *Let $f(r, t)$ and $rf_t(r, t) \in C(\overline{D})$. Then the generalized solution $u(r, t)$ of Problem $P21$ belongs to $C^2(\overline{D} \setminus (0, 0))$ and has the following asymptotic expansion at $(0, 0)$*

$$u(r, t) = \sum_{k=0}^{\lfloor n/2 \rfloor} r^{-1} (r+t)^{-(n-2k)} \beta_k^n F_k^n(r, t) + F^n(r, t),$$

where:

- 1) the functions $F_k^n \in C^2(\overline{D} \setminus (0, 0))$ are independent of f , bounded, and satisfy $F_k^n(t, t) \equiv \text{const} \neq 0$.
- 2) the function $F^n \in C^2(\overline{D} \setminus (0, 0))$ satisfies the following estimates:

$$\begin{aligned} |F^0(r, t)| &\leq C \|rf\|_{C(D)}, \\ |(F^0)_r(r, t)| + |(F^0)_t(r, t)| &\leq Cr^{-1} \|rf\|_{C(D)}, \end{aligned}$$

and for $n \geq 1$

$$(6.2) \quad |F^n(r, t)| \leq CA_1 n^2 (1 + |\ln r|).$$

If also rf and $rf_t \in C^1(D)$, then

$$(6.3) \quad |F^n(r, t)| \leq Cr^{-1}(A_0n^2 + A_2),$$

and if additionally $rf_{tt} \in C^1(D)$ and $r^2f_{trr} \in C(D)$, then

$$(6.4) \quad |(F^n)_r(r, t)| + |(F^n)_t(r, t)| \leq Cr^{-3}(A_1n^2 + CA_3),$$

where in all inequalities the constant C is independent of n and f .

The proof of this result is quite long and technical and we leave it for the last Section 7. Here we will use Theorem 6.1 to prove Theorem 1.2 and Theorem 1.3.

Proof of Theorem 1.2. Assume that the right-hand side function f is a harmonic polynomial (1.4). Then the unique generalized solution $u(x, t)$ also is a harmonic polynomial (5.1), according to Theorem 5.2. Furthermore, the functions $u_n^m(r, t)$ are solutions of Problem $P21$ with right-hand side f_n^m that can be represented as

$$(6.5) \quad f_n^m(r, t) := \int_0^\pi \left(\int_0^{2\pi} f(r, \theta, \varphi, t) Y_n^m(\theta, \varphi) d\varphi \right) \sin \theta d\theta.$$

When $f \in C^1(\overline{\Omega})$ we have $f_n^m \in C^1(\overline{D})$ and obviously

$$(6.6) \quad \|f_n^m(r, t)\|_{C^1(D)} \leq C \|f(x, t)\|_{C^1(\Omega)}.$$

The definition of functions $W_{k,m}^n$ from Lemma 1.1 and (6.5) give the identity

$$(6.7) \quad \int_0^{\frac{1}{2}} \left(\int_t^{1-t} E_k^n(r, t) f_n^m(r, t) r^2 dr \right) dt = \int_\Omega W_{k,m}^n(x, t) f(x, t) dx dt,$$

which shows that $\beta_k^n = \beta_{k,m}^n$, according to their definitions (6.1) and (1.5).

Now we can apply Theorem 6.1 for the functions $u_n^m(r, t)$ and $f_n^m(r, t)$. Using (6.6) and (6.7) we get the expansion

$$u_n^m(r, t) = \sum_{k=0}^{[n/2]} r^{-1}(r+t)^{-(n-2k)} \beta_{k,m}^n F_k^{n,m}(r, t) + F^{n,m}(r, t),$$

where $|F^{n,m}(r, t)| \leq C \|f\|_{C^1(\Omega)}$, $|F_k^{n,m}(r, t)| \leq C$ and $F_k^{n,m}(t, t) = \text{const} \neq 0$. Summing up over n and m one gets the desired expansion.

Finally, to prove property (iii), let us fix a direction $(\alpha, 1) := (\alpha_1, \alpha_2, \alpha_3, 1)$ with $\alpha_1 = \sin \theta_0 \cos \varphi_0$, $\alpha_2 = \sin \theta_0 \sin \varphi_0$ and $\alpha_3 = \cos \theta_0$. Then for the functions $F_{k,m}^n$ from (1.6) we have

$$F_{k,m}^n(\alpha t, t) := 2^{(2k-n+1)/2} F_k^{n,m}(t, t) Y_n^m(\varphi_0, \theta_0)$$

and thus, there are some nonzero constants $C_{p,k,m}$ such that

$$\lim_{t \rightarrow +0} F_p(\alpha t, t) = \sum_{k=0}^{[(l-p+1)/2]} \sum_{m=1}^{2p+4k-1} C_{p,k,m} \beta_{k,m}^{p+2k-1} Y_{p+2k-1}^m(\varphi_0, \theta_0).$$

Therefore the property (iii) follows from the fact that the spherical functions Y_n^m are linearly independent. \square

Besides Theorem 6.1, the proof of Theorem 1.3 relies on and the estimates from Section 3 for the special functions.

Proof of Theorem 1.3. The function $f \in C^6(\bar{\Omega})$ can be represented as

$$f(x, t) = \sum_{n=0}^{\infty} \sum_{m=1}^{2n+1} f_n^m(|x|, t) Y_n^m(x).$$

The generalized solution of Problem *P2* could be formally written in the form

$$(6.8) \quad u(x, t) = \sum_{n=0}^{\infty} \sum_{m=1}^{2n+1} u_n^m(|x|, t) Y_n^m(x),$$

where $u_n^m(r, t)$ is the solution of Problem *P21* with right-hand side f_n^m . We will prove that the series (6.8) and its derivatives are uniformly convergent in $\bar{\Omega} \cap \{r \geq \varepsilon\}$ for $\varepsilon > 0$, and that u is bounded.

According to Lemma 3.3 the series for f and its first derivatives uniformly converge. For the derivatives with respect to x_i and with respect to r , there holds the relation

$$\sum_{i=1}^3 x_i \frac{\partial}{\partial x_i} f(x, t) = \sum_{n=0}^{\infty} \sum_{m=1}^{2n+1} r \frac{\partial}{\partial r} f_n^m(r, t) Y_n^m(x),$$

and therefore $r(f_n^m)_r \in C(\bar{D})$. A similar argument shows that $r f_n^m$, $r(f_n^m)_t$, $r(f_n^m)_{tt} \in C^1(\bar{D})$ and $r^2(f_n^m)_{trr} \in C(\bar{D})$ and we can apply Theorem 6.1.

First, using to Lemma 3.3 case 1) with $k = 3$ and $k = 2$ we see that for $n > 0$

$$(6.9) \quad \|f_n^m\|_{C(D)} \leq C n^{-6} \|f\|_{C^6(\bar{\Omega})},$$

$$(6.10) \quad \|r f_n^m\|_{C^1(D)} + \|r(f_n^m)_t\|_{C^1(D)} \leq C n^{-4} \|f\|_{C^6(\bar{\Omega})}.$$

On the other hand, Lemma 3.3 case 2) with $k = 2$ and $k = 1$ gives

$$(6.11) \quad \sum_{m=1}^{2n+1} \|f_n^m\|_{C(D)} \leq C n^{-4} r \|f\|_{C^5(\bar{\Omega})},$$

$$(6.12) \quad \sum_{m=1}^{2n+1} \|(f_n^m)_t\|_{C(D)} \leq C n^{-4} r \|f_t\|_{C^5(\bar{\Omega})},$$

$$(6.13) \quad \sum_{m=1}^{2n+1} (\|r f_n^m\|_{C^1(D)} + \|r f_n^m\|_{C^1(D)} + \|r(f_n^m)_{tt}\|_{C^1(D)} + \|r^2(f_n^m)_{trr}\|_{C(D)}) \leq C n^{-2} r \|f\|_{C^6(\bar{\Omega})}.$$

Next we study the series (6.8). Using the notations of Theorem 6.1, we know that when $\beta_k^n = 0$ for $k = 0, \dots, [n/2]$, the function $u_n^m(r, t)$ satisfies

$$u_n^m(r, t) = F^n(r, t)$$

and then for $n > 0$ from (6.2), (6.3) and (6.4) we have estimates:

$$(6.14) \quad |u_n^m(r, t)| \leq Cn^2 (\|f_n^m\|_{C(D)} + \|r(f_n^m)_t\|_{C(D)}) (1 + |\ln r|)$$

or alternatively

$$(6.15) \quad |u_n^m(r, t)| \leq Cr^{-1} (n^2 \|f_n^m\|_{C(D)} + \|rf_n^m\|_{C^1(D)} + \|r(f_n^m)_t\|_{C^1(D)}),$$

and for the derivatives

$$(6.16) \quad \begin{aligned} & |(u_n^m)_r(r, t)| + |(u_n^m)_t(r, t)| \\ & \leq Cr^{-3} \left(n^2 \|f_n^m\|_{C(D)} + n^2 \|r(f_n^m)_t\|_{C(D)} \right. \\ & \quad \left. + \|rf_n^m\|_{C^1(D)} + \|rf_n^m\|_{C^1(D)} + \|r(f_n^m)_{tt}\|_{C^1(D)} + \|r^2(f_n^m)_{trr}\|_{C(D)} \right). \end{aligned}$$

Then applying the estimate (3.9) for the spherical functions from Corollary 3.1 and substituting (6.11) and (6.12) in (6.14) we find

$$\begin{aligned} \sum_{m=1}^{2n+1} |u_n^m Y_n^m| & \leq Cn^{-5/2} (1 + |\ln r|) \sum_{m=1}^{2n+1} (\|f_n^m\|_{C(D)} + \|r(f_n^m)_t\|_{C(D)}) \\ & \leq Cn^{-3/2} (\|f\|_{C^5(\bar{\Omega})} + \|f_t\|_{C^5(\bar{\Omega})}). \end{aligned}$$

On the other hand, first using (6.15) with (6.9) and (6.10), and then (3.12) for the sum of derivatives of Y_n^m , we get

$$\begin{aligned} & \sum_{m=1}^{2n+1} \sum_{|\alpha|=1} |u_n^m D_{(x,t)}^\alpha Y_n^m| \\ & \leq Cn^{-4} r^{-1} \|f\|_{C^6(\bar{\Omega})} \sum_{m=1}^{2n+1} \sum_{|\alpha|=1} |D_{(x,t)}^\alpha Y_n^m| \leq Cn^{-2} r^{-2} \|f\|_{C^6(\bar{\Omega})}. \end{aligned}$$

Combining (6.16) for the derivatives of u with (6.11), (6.12) and (6.13) gives

$$\begin{aligned} & \sum_{m=1}^{2n+1} \sum_{|\alpha|=1} |Y_n^m D_{(x,t)}^\alpha u_n^m| \\ & \leq Cn^{5/2} r^{-3} \sum_{m=1}^{2n+1} (\|f_n^m\|_{C(D)} + \|r(f_n^m)_t\|_{C(D)}) \\ & \quad + Cn^{1/2} r^{-3} \sum_{m=1}^{2n+1} (\|rf_n^m\|_{C^1(D)} + \|rf_n^m\|_{C^1(D)} + \|r(f_n^m)_{tt}\|_{C^1(D)} + \|r^2(f_n^m)_{trr}\|_{C(D)}) \\ & \leq Cn^{-3/2} r^{-2} \|f\|_{C^6(\bar{\Omega})}. \end{aligned}$$

For the case $n = 0$ we have $Y_0^1 = \text{const}$ and representation (6.5) shows that $|f_0^1| \leq C\|f\|_{C(\bar{\Omega})}$. Therefore Theorem 6.1 gives

$$|u_0^1 Y_0^1| \leq C\|f\|_{C(\bar{\Omega})},$$

$$\sum_{|\alpha|=1} |D_{(x,t)}^\alpha u_0^1 Y_0^1| \leq Cr^{-1} \|f\|_{C(\bar{\Omega})}.$$

After this preparation, we are ready to estimate the Fourier series (6.8) and its first derivatives:

$$\begin{aligned} |u(x, t)| &\leq C\|f\|_{C(\bar{\Omega})} + C\left(\|f\|_{C^5(\bar{\Omega})} + \|f_t\|_{C^5(\bar{\Omega})}\right) \sum_{n=1}^{\infty} n^{-2} \\ &\leq C\left(\|f\|_{C^5(\bar{\Omega})} + \|f_t\|_{C^5(\bar{\Omega})}\right), \end{aligned}$$

and

$$\begin{aligned} &\sum_{|\alpha|=1} |D_{(x,t)}^\alpha u(x, t)| \\ &\leq Cr^{-1}\|f\|_{C(\bar{\Omega})} + Cr^{-2}\|f\|_{C^6(\bar{\Omega})} \sum_{n=1}^{\infty} n^{-3/2} + Cr^{-2}\|f\|_{C^6(\bar{\Omega})} \sum_{n=1}^{\infty} n^{-2} \\ &\leq Cr^{-2}\|f\|_{C^6(\bar{\Omega})}. \end{aligned}$$

Therefore we have $u \in C^1(\bar{\Omega} \setminus O)$ since for each fixed $\varepsilon > 0$ the series (6.8) uniformly converges in the set $\bar{\Omega} \cap \{r \geq \varepsilon\}$ and the same holds for its first derivatives.

Finally, we will prove that the function $u(x, t)$ defined as the series (6.8) is the generalized solution of Problem *P2*. First, notice the function $u_n^m(|x|, t)Y_n^m(x)$ is the generalized solution of Protter Problem *P2* with right hand side function $f_n^m(|x|, t)Y_n^m(x)$. Thus $u \in C^1(\bar{\Omega} \setminus O)$ satisfies the boundary conditions $u_t|_{\Sigma_0} = 0$ and $u|_{\Sigma_1} = 0$ just like all the terms $u_n^m Y_n^m$. The proof of the case 2) from the Definition 1.1 is straightforward – for a test function w and $l \in \mathbb{N}$ we have

$$\begin{aligned} &\int_{\Omega} \sum_{n=0}^l \sum_{m=1}^{2n+1} \{(u_n^m Y_n^m)_t w_t - (u_n^m Y_n^m)_{x_1} w_{x_1} \\ &\quad - (u_n^m Y_n^m)_{x_2} w_{x_2} - (u_n^m Y_n^m)_{x_3} w_{x_3} - f_n^m Y_n^m w\} dx dt = 0 \end{aligned}$$

and the uniform convergence in $\Omega \cap \text{supp}(w)$ of the series (6.8) and its first derivatives, allow us to take the limit $l \rightarrow \infty$ in this equality. Therefore

$$\int_{\Omega} \{u_t w_t - u_{x_1} w_{x_1} - u_{x_2} w_{x_2} - u_{x_3} w_{x_3} - f w\} dx dt = 0$$

and we see that $u(x, t)$ is the generalized solution of Problem *P2* with right-hand side $f(x, t)$. \square

7. THE PROOF OF THE ASYMPTOTIC EXPANSION IN THE TWO-DIMENSIONAL CASE

The proof of Theorem 6.1 is based on the results stated in Section 5. In particular, according to Lemma 5.1 the solution $u(r, t)$ of Problem *P21* can be constructed with help of the substitutions $\xi = (r + t)/2$, $\eta = (r - t)/2$ as

$$u(\xi + \eta, \xi - \eta) = (\xi + \eta)^{-1} v(\xi, \eta),$$

where $v(\xi, \eta)$ is defined by the formulas (5.4), (5.5), (5.6) and

$$g(\xi, \eta) = (\xi + \eta)f(\xi + \eta, \xi - \eta).$$

One can see that generally the integral in (5.5) blows up when ξ approaches 0, and thus $v(\xi, \eta)$ has singularity at $(0, 0)$ even for smooth functions $g(\xi, \eta)$.

Proof of Theorem 6.1:

A. Proof of the asymptotic formula.

We will study the behavior of the function

$$v(\xi, \eta) = (\xi + \eta)u(\xi + \eta, \xi - \eta),$$

given by the integral representation (5.4) from Lemma 5.1. The smoothness of $v(\xi, \eta)$ in (5.4) away from the point $(0, 0)$ follows directly from the smoothness of the function $G(\xi)$. Next we will derive the asymptotic expansion of $v(\xi, \eta)$ at $(0, 0)$.

First we will find the relation between the constants β_k^n and the function $G(\xi)$ defined by (5.6). Let us compute the integral

$$\begin{aligned} \int_0^{\frac{1}{2}} \xi^k G(\xi) d\xi &= \int_0^{\frac{1}{2}} \int_{\xi}^{\frac{1}{2}} \int_0^{\xi} \xi^k P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 d\xi_1 d\xi \\ &\quad - \int_0^{\frac{1}{2}} \int_0^{\xi} \xi^k P_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) g \left(\frac{1}{2}, \eta_1 \right) d\eta_1 d\xi - \int_0^{\frac{1}{2}} \int_{\xi}^{\frac{1}{2}} \xi^k P_n \left(\frac{\xi}{\xi_1} \right) g(\xi_1, 0) d\xi_1 d\xi \\ &= \int_0^{\frac{1}{2}} \int_0^{\xi_1} \left(\int_{\eta_1}^{\xi_1} \xi^k P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) d\xi \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 d\xi_1 \\ &\quad - \int_0^{\frac{1}{2}} \left(\int_{\eta_1}^{\frac{1}{2}} \xi^k P_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) d\xi \right) g \left(\frac{1}{2}, \eta_1 \right) d\eta_1 - \int_0^{\frac{1}{2}} \left(\int_0^{\xi_1} \xi^k P_n \left(\frac{\xi}{\xi_1} \right) d\xi \right) g(\xi_1, 0) d\xi_1 \\ &= - \int_0^{\frac{1}{2}} \int_0^{\xi_1} \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) \left(\int_{\eta_1}^{\xi_1} \xi^k P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) d\xi \right) g(\xi_1, \eta_1) d\eta_1 d\xi_1 \\ &= - \int_0^{\frac{1}{2}} \int_0^{\xi_1} \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) h_k(\xi_1, \eta_1) g(\xi_1, \eta_1) d\eta_1 d\xi_1. \end{aligned}$$

Now we can apply Lemma 5.2 to get

$$\left(\frac{\partial}{\partial \xi} - \frac{\partial}{\partial \eta} \right) h_{n-2i}(\xi, \eta) \Big|_{\substack{\xi = (r+t)/2 \\ \eta = (r-t)/2}} = c_i^n r E_i^n(r, t)$$

and we conclude that

$$(7.1) \quad \begin{aligned} \int_0^{\frac{1}{2}} \xi^{n-2i} G(\xi) d\xi &= - \int_0^{\frac{1}{2}} \left(\int_0^{\xi} \left(\frac{\partial}{\partial \xi} - \frac{\partial}{\partial \eta} \right) h_{n-2i}^n(\xi, \eta) g(\xi, \eta) d\eta \right) d\xi \\ &= -c_i^n \int_0^{\frac{1}{2}} \left(\int_t^{1-t} E_i^n(r, t) f(r, t) r^2 dr \right) dt = -c_i^n \beta_i^n. \end{aligned}$$

Next we consider the case $n \geq 1$. The simpler case $n = 0$ will be discussed separately later in the proof.

Let us expand the Legendre polynomial P_n in formula (5.5) using (3.2):

$$P_n(x) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} x^{n-2k},$$

where $a_{2k} \neq 0$. For $\tau(\xi)$ we get

$$\tau(\xi) = \int_{\frac{1}{2}}^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) G(\xi_1) d\xi_1 = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} \xi^{-n+2k} \int_{\frac{1}{2}}^{\xi} \xi_1^{n-2k} G(\xi_1) d\xi_1.$$

Applying (7.1) we find

$$(7.2) \quad \tau(\xi) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} \xi^{-n+2k} \left(c_i^n \beta_i^n + \int_0^{\xi} \xi_1^{n-2k} G(\xi_1) d\xi_1 \right) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} c_i^n \beta_i^n \xi^{-n+2k} + \psi(\xi),$$

where

$$(7.3) \quad \psi(\xi) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} \xi^{-n+2k} \int_0^{\xi} \xi_1^{n-2k} G(\xi_1) d\xi_1 = \int_0^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) G(\xi_1) d\xi_1.$$

Now, we want to estimate the function $\psi(\xi)$. First, let us look more carefully to the representation (5.6) of the function $G(\xi)$ with $g = r f(r, t)$:

$$(7.4) \quad \begin{aligned} G(\xi) &= \int_{\xi}^{\frac{1}{2}} \left(\int_0^{\xi} P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1 \\ &\quad - \int_0^{\xi} P_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) g \left(\frac{1}{2}, \eta_1 \right) d\eta_1 - \int_{\xi}^{\frac{1}{2}} P_n \left(\frac{\xi}{\xi_1} \right) g(\xi_1, 0) d\xi_1. \end{aligned}$$

Since all the arguments of P_n here are in the interval $[0, 1]$ and $|P_n(x)| \leq 1$ for $x \in [0, 1]$, it is obvious that

$$(7.5) \quad |G(\xi)| \leq CA_1$$

and therefore

$$(7.6) \quad |\psi(\xi)| \leq CA_1\xi,$$

where the constant C is independent of n and f .

We will need also a more accurate estimate for $\psi(\xi)$ with higher power of ξ . The first two integrals in (7.4) are bounded by $2A_1\xi$. For the last term

$$J_n(\xi) := \int_{\xi}^{\frac{1}{2}} P_n\left(\frac{\xi}{\xi_1}\right) g(\xi_1, 0) d\xi_1$$

more computations are required. First, using the estimate (3.5) from Lemma 3.1, we have

$$\begin{aligned} |J_n(\xi) - J_n(0)| &= \left| \int_{\xi}^{\frac{1}{2}} P_n\left(\frac{\xi}{\xi_1}\right) g(\xi_1, 0) d\xi_1 - P_n(0) \int_0^{\frac{1}{2}} g(\xi_1, 0) d\xi_1 \right| \\ &\leq n \int_{\xi}^{\frac{1}{2}} \frac{\xi}{\xi_1} |g(\xi_1, 0)| d\xi_1 + \left| P_n(0) \int_0^{\xi} g(\xi_1, 0) d\xi_1 \right| \leq CA_1 n \xi^{\varepsilon}, \end{aligned}$$

with some $\varepsilon \in (0, 1)$ and the constant C is independent of n . Therefore

$$(7.7) \quad |G(\xi) + J_n(0)| \leq CA_1 n \xi^{\varepsilon}.$$

Now, notice that the value of $\psi(\xi)$ will not change if we add $J_n(0)$ to $G(\xi)$. This is based on the equality

$$\int_0^{\xi} P_n\left(\frac{\xi_1}{\xi}\right) J_n(0) d\xi_1 \equiv P_n(0) \int_0^{\frac{1}{2}} g(\xi_1, 0) d\xi_1 \int_0^{\xi} P_n\left(\frac{\xi_1}{\xi}\right) d\xi_1 = 0,$$

that holds, because when n is odd number $P_n(0) = 0$, while for even indices n , $n > 0$, the polynomial P_n is an even function and by the definition of the Legendre polynomials (see (3.2)) follows

$$\int_0^{\xi} P_n\left(\frac{\xi_1}{\xi}\right) d\xi_1 = \xi \int_0^1 P_n(t) dt = \frac{1}{2} \xi \int_{-1}^1 P_n(t) dt = c\xi \int_{-1}^1 \frac{d^n}{dt^n} \{(1-t^2)^n\} dt = 0.$$

Thus

$$\psi(\xi) = \psi(\xi) + \int_0^{\xi} P_n\left(\frac{\xi_1}{\xi}\right) J_n(0) d\xi_1 = \int_0^{\xi} P_n\left(\frac{\xi_1}{\xi}\right) [G(\xi_1) + J_n(0)] d\xi_1$$

and we can apply (7.7) to conclude that for $n > 0$ there is $\varepsilon > 0$ and a constant C independent of n and f , such that

$$(7.8) \quad |\psi(\xi)| \leq A_1 C n \xi^{1+\varepsilon}.$$

Now we apply the expansion (7.2) of $\tau(\xi)$ in the definition (5.4) of $v(\xi, \eta)$ and find that

$$\begin{aligned} v(\xi, \eta) &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} c_k^n \beta_k^n \xi^{-n+2k} + \psi(\xi) \\ &+ \int_{\xi}^{\frac{1}{2}} \tau(\xi_1) \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 + F_1(\xi, \eta), \end{aligned}$$

where the function F_1 is smooth (see (5.4)) and $|F_1(\xi, \eta)| \leq A_1 C\xi$. Consider the term

$$\begin{aligned} &\int_{\xi}^{\frac{1}{2}} \tau(\xi_1) \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} c_k^n \beta_k^n \int_{\xi}^{\frac{1}{2}} \xi_1^{-n+2k} \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 \\ &+ \int_{\xi}^{\frac{1}{2}} \psi(\xi_1) \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1. \end{aligned}$$

According to (7.8) the last integral is bounded:

$$\left| \int_{\xi}^{\frac{1}{2}} \xi_1^{1+\varepsilon} \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 \right| \leq \frac{C\xi\eta}{\xi + \eta} \int_{\xi}^{\frac{1}{2}} \xi_1^{\varepsilon-1} d\xi_1 \leq C'\xi.$$

Thus, we find the expansion

$$(7.9) \quad v(\xi, \eta) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} a_{2k} c_k^n \beta_k^n \xi^{-n+2k} G_k^n(\xi, \eta) + \psi_1(\xi, \eta),$$

where

$$G_k^n(\xi, \eta) := 1 + \xi^{n-2k} \int_{\xi}^{\frac{1}{2}} \xi_1^{-n+2k} \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1$$

and $|\psi_1(\xi, \eta)| \leq A_1 C\xi$. Let us point out that $G_k^n(\xi, 0) = 1$, while the fact that the functions G_k^n are bounded, follows from the estimate

$$\left| \int_{\xi}^{\frac{1}{2}} \xi_1^{-n+2k} \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 \right| \leq C\xi^{-n+2k}.$$

The representation (7.9) holds for $n > 0$, while when $n = 0$ we have simply $P_0 \equiv 1$ and can substitute it in formulas (5.4), (5.5) and (5.6) for the solution v of Problem *P22*. Straightforward computations lead to

$$(7.10) \quad v(\xi, \eta) = a_0 c_0^0 \beta_0^0 + \int_0^\xi G(\xi_1) d\xi_1 - \int_\xi^{\frac{1}{2}} \left(\int_0^\eta g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1,$$

where

$$G(\xi) := - \int_0^\xi g(\xi, \eta_1) d\eta_1 - \int_\xi^{\frac{1}{2}} g(\xi_1, \xi) d\xi_1.$$

Therefore, the representation (7.9) obviously stays true in the case $n = 0$.

Finally, let us return to the generalized solution $u(r, t)$ of Problem *P21* and to the coordinates r and t :

$$u(r, t) = r^{-1} v \left(\frac{r+t}{2}, \frac{r-t}{2} \right) = r^{-1} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \beta_k^n (r+t)^{-n+2k} F_k^n(r, t) + F^n(r, t).$$

Here, the function $F^n(r, t)$ is given by

$$(7.11) \quad F^n(r, t) := r^{-1} \psi_1 \left(\frac{r+t}{2}, \frac{r-t}{2} \right)$$

and therefore $|F^n(r, t)| \leq CA_1$, while functions $F_k^n(r, t)$ defined by

$$F_k^n(r, t) := 2^{n-2k} a_{2k} c_k^n G_k^n \left(\frac{r+t}{2}, \frac{r-t}{2} \right)$$

are independent on f and are obviously bounded. To complete the proof of the case 1), notice that this definition gives

$$F_k^n(t, t) = 2^{n-2k} a_{2k} c_k^n \neq 0.$$

B. Proof of the estimates of F^n .

Next, for $n \geq 1$, we will estimate the function F^n and its first derivatives.

First, we will study the behavior of F^n at $(0, 0)$. The function $F^n(r, t)$ is given by (7.11), where $\psi_1(\xi, \eta)$ is defined in (7.9) as

$$(7.12) \quad \psi_1(\xi, \eta) = \psi(\xi) + \int_\xi^{\frac{1}{2}} \psi(\xi_1) \frac{\partial}{\partial \xi_1} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) d\xi_1 + F_1(\xi, \eta),$$

and

$$(7.13) \quad F_1(\xi, \eta) = - \int_\xi^{\frac{1}{2}} \left(\int_0^\eta P_n \left(\frac{(\xi - \eta)(\xi_1 - \eta_1) + 2\xi_1\eta_1 + 2\xi\eta}{(\xi_1 + \eta_1)(\xi + \eta)} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1.$$

Since the argument of the Legendre polynomial P_n in (7.13) varies in the interval $[0, 1]$

$$|F_1(\xi, \eta)| \leq CA_0\xi,$$

where the constant C is independent of n and f . Thus applying the estimate (7.6) for ψ we get

$$|\psi_1(\xi, \eta)| \leq CA_1\xi + CA_1n^2 \int_{\xi}^{\frac{1}{2}} \frac{\xi\eta}{\xi_1(\xi + \eta)} d\xi_1 + CA_1\xi \leq C_1A_1n^2\xi |\ln \xi|.$$

Therefore

$$|F^n(r, t)| \leq C_2A_1n^2 (1 + |\ln r|),$$

where the constant C_2 is independent of n and f , i.e., (6.2) holds.

Now we consider the derivatives of F^n . Thus we need to evaluate the derivatives of $\psi_1(\xi, \eta)$ – integrating (7.12) by parts we get
(7.14)

$$\psi_1(\xi, \eta) = P_n \left(\frac{\xi - \eta + 4\xi\eta}{\xi + \eta} \right) \psi(1/2) - \int_{\xi}^{\frac{1}{2}} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) \psi'(\xi_1) d\xi_1 + F_1(\xi, \eta).$$

For $\psi'(\xi)$ defined by (7.3), using (7.5) we find

$$|\psi'(\xi)| = \left| G(\xi) - \int_0^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) \frac{\xi_1}{\xi^2} G(\xi_1) d\xi_1 \right| \leq Cn^2 \max |G(\xi)| \leq CA_1n^2,$$

where the constant C is independent of n and f . In fact, we can remove the coefficient n^2 here for smoother functions g . In order to do this, let us rewrite ψ' as

$$(7.15) \quad \psi'(\xi) = \xi^{-1} \int_0^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) \frac{\partial}{\partial \xi_1} \{ \xi_1 G(\xi_1) \} d\xi_1.$$

For $(\psi_1)_{\xi}$ and $(\psi_1)_{\eta}$ we will need also ψ'' :

$$(7.16) \quad \psi''(\xi) = \xi^{-2} \int_0^{\xi} P_n \left(\frac{\xi_1}{\xi} \right) \frac{\partial^2}{\partial \xi_1^2} \{ \xi_1 G(\xi_1) \} d\xi_1.$$

Then estimates for the derivatives of $G(\xi)$ are required:

$$\begin{aligned}
G'(\xi) &= - \int_0^\xi \frac{\partial}{\partial \xi} g(\xi, \eta_1) d\eta_1 - \int_\xi^{\frac{1}{2}} \frac{\partial}{\partial \xi} g(\xi_1, \xi) d\xi_1 \\
&\quad - \int_0^\xi P'_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) \frac{2\xi^2 - \eta_1}{\xi^2(2\eta_1 + 1)} g \left(\frac{1}{2}, \eta_1 \right) d\eta_1 - \int_\xi^{\frac{1}{2}} P'_n \left(\frac{\xi}{\xi_1} \right) \frac{g(\xi_1, 0)}{\xi_1} d\xi_1 \\
&\quad + \int_\xi^{\frac{1}{2}} \left(\int_0^\xi \frac{\partial}{\partial \xi} P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1.
\end{aligned}$$

Notice that for the argument of P_n in the last term

$$(7.17) \quad \left(\xi \frac{\partial}{\partial \xi} + \xi_1 \frac{\partial}{\partial \xi_1} + \eta_1 \frac{\partial}{\partial \eta_1} \right) \frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} = 0$$

since the function is homogenous. Therefore we can replace there the derivative with respect to ξ with $[-\xi_1 \partial / \partial \xi_1 - \eta_1 \partial / \partial \eta_1] / \xi$. Integrating by parts we find

$$\begin{aligned}
G'(\xi) &= -g(1/2, \xi) - \frac{1}{2\xi} \int_0^\xi P_n \left(\frac{\eta_1 + 2\xi^2}{\xi(1 + 2\eta_1)} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) \Big|_{\xi_1=1/2} d\eta_1 \\
&\quad + g(\xi, 0) - \int_\xi^{\frac{1}{2}} P'_n \left(\frac{\xi}{\xi_1} \right) \frac{g(\xi_1, 0)}{\xi_1} d\xi_1 - \int_0^\xi P'_n \left(\frac{\eta_1 + 2\xi^2}{\xi(2\eta_1 + 1)} \right) \frac{(2\xi^2 - \eta_1)g(1/2, \eta_1)}{\xi^2(2\eta_1 + 1)} d\eta_1 \\
&\quad + \frac{1}{\xi} \int_\xi^{\frac{1}{2}} \left(\int_0^\xi P_n \left(\frac{\xi_1 \eta_1 + \xi^2}{\xi(\xi_1 + \eta_1)} \right) \left(2 + \xi_1 \frac{\partial}{\partial \xi_1} + \eta_1 \frac{\partial}{\partial \eta_1} \right) \left(\frac{\partial}{\partial \xi_1} - \frac{\partial}{\partial \eta_1} \right) g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1
\end{aligned}$$

Recall that $g(\xi, \eta) = rf(r, t)$, and thus $g(\xi, 0)/\xi = f(r, r)$ and in the last integral we have

$$\left(\xi \frac{\partial}{\partial \xi} + \eta \frac{\partial}{\partial \eta} \right) \left(\frac{\partial}{\partial \xi} - \frac{\partial}{\partial \eta} \right) g(\xi, \eta) \Big|_{\substack{\xi=(r+t)/2 \\ \eta=(r-t)/2}} = \left(r \frac{\partial}{\partial r} + t \frac{\partial}{\partial t} \right) rf_t(r, t).$$

Hence, using (3.4) from Lemma 3.1, we get that

$$(7.18) \quad |G'(\xi)| \leq CA_0 n^2 + CA_2.$$

Analogously for $G''(\xi)$, differentiating one more time the expression for $G'(\xi)$, applying again (7.17) in the last integral, and integrating by parts the last three terms, we find the estimate

$$(7.19) \quad |G''(\xi)| \leq CA_1 n^2 \xi^{-1} (1 - 2\xi)^{-1} + CA_3 \xi^{-1}.$$

Applying (7.5), (7.18), (7.19) to ψ and ψ' in (7.15), (7.16) we find

$$(7.20) \quad |\psi'(\xi)| \leq CA_0 n^2 + CA_2,$$

and

$$(7.21) \quad |\psi''(\xi)| \leq CA_1 n^2 \xi^{-2} |\ln(1 - 2\xi)| + CA_3 \xi^{-1}.$$

To evaluate the derivatives of ψ_1 we need also to study the derivatives of F_1 defined by (7.13):

$$\begin{aligned} (F_1)_\eta &= - \int_{\xi}^{\frac{1}{2}} g(\xi_1, \eta) d\xi_1 \\ &+ \int_{\xi}^{\frac{1}{2}} \left(\int_0^{\eta} P'_n \left(\frac{(\xi - \eta)(\xi_1 - \eta_1) + 2\xi_1 \eta_1 + 2\xi \eta}{(\xi_1 + \eta_1)(\xi + \eta)} \right) \frac{2(\xi_1 - \xi)(\xi + \eta_1)}{(\xi_1 + \eta_1)(\xi + \eta)^2} g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1; \\ (F_1)_\xi &= \int_0^{\eta} g(\xi, \eta_1) d\eta_1 \\ &+ \int_{\xi}^{\frac{1}{2}} \left(\int_0^{\eta} P'_n \left(\frac{(\xi - \eta)(\xi_1 - \eta_1) + 2\xi_1 \eta_1 + 2\xi \eta}{(\xi_1 + \eta_1)(\xi + \eta)} \right) \frac{2(\xi_1 + \eta)(\eta - \eta_1)}{(\xi_1 + \eta_1)(\xi + \eta)^2} g(\xi_1, \eta_1) d\eta_1 \right) d\xi_1. \end{aligned}$$

Notice that in these integrals $0 \leq \eta_1 \leq \eta \leq \xi \leq \xi_1$ and therefore

$$\frac{(\xi_1 - \xi)(\xi + \eta_1)}{(\xi_1 + \eta_1)(\xi + \eta)^2} \leq \frac{1}{\xi + \eta} \quad \text{and} \quad \frac{(\xi_1 + \eta)(\eta - \eta_1)}{(\xi_1 + \eta_1)(\xi + \eta)^2} \leq \frac{1}{\xi + \eta}.$$

Then it follows from Lemma 3.1 that

$$|(F_1)_\xi(\xi, \eta)| + |(F_1)_\eta(\xi, \eta)| \leq CA_1 n^2.$$

Now we are ready to estimate the derivatives of $\psi(\xi, \eta)$ from (7.14). For $(\psi_1)_\eta(\xi, \eta)$ after integration by part we have

$$\begin{aligned} (\psi_1)_\eta &= - \int_{\xi}^{\frac{1}{2}} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) \frac{\partial}{\partial \xi_1} \left[\frac{\xi_1(\xi - \xi_1)}{\eta(\xi + \eta)} \psi'(\xi_1) \right] d\xi_1 + (F_1)_\eta \\ &+ P'_n \left(\frac{\xi - \eta + 4\xi\eta}{\xi + \eta} \right) \frac{2\xi(2\xi - 1)}{(\xi + \eta)^2} \psi(1/2) + P_n \left(\frac{\xi - \eta + 4\xi\eta}{\xi + \eta} \right) \frac{2\xi - 1}{4\eta(\xi + \eta)} \psi'(1/2). \end{aligned}$$

Then, applying (7.20) and (7.21), it follows

$$|(\psi_1)_\eta| \leq C(\xi + \eta)^{-2} [A_1 n^2 + CA_3].$$

Similarly, for $(\psi_1)_\xi$ we find

$$\begin{aligned} (\psi_1)_\xi &= - \int_{\xi}^{\frac{1}{2}} P_n \left(\frac{(\xi - \eta)\xi_1 + 2\xi\eta}{\xi_1(\xi + \eta)} \right) \frac{\partial}{\partial \xi_1} \left[\frac{\xi_1(\eta + \xi_1)}{\xi(\xi + \eta)} \psi'(\xi_1) \right] d\xi_1 + (F_1)_\xi \\ &+ P'_n \left(\frac{\xi - \eta + 4\xi\eta}{\xi + \eta} \right) \frac{2\eta(2\eta + 1)}{\xi + \eta} \psi(1/2) + P_n \left(\frac{\xi - \eta + 4\xi\eta}{\xi + \eta} \right) \frac{2\eta + 1}{4\xi(\xi + \eta)} \psi'(1/2), \end{aligned}$$

and thus we get

$$|(\psi_1)_\xi| \leq C(\xi + \eta)^{-2} (A_1 n^2 + CA_3).$$

Finally, to prove (6.4) notice that

$$(F^n)_r(r, t) = -r^{-2}\psi_1(\xi, \eta) \Big|_{\substack{\xi=(r+t)/2; \\ \eta=(r-t)/2}} + \frac{1}{2}r^{-1}((\psi_1)_\xi + (\psi_1)_\eta) \Big|_{\substack{\xi=(r+t)/2; \\ \eta=(r-t)/2}}$$

and

$$(F^n)_t(r, t) = \frac{1}{2}r^{-1}((\psi_1)_\xi - (\psi_1)_\eta) \Big|_{\substack{\xi=(r+t)/2; \\ \eta=(r-t)/2}}$$

and therefore

$$|(F^n)_r(r, t)| + |(F^n)_t(r, t)| \leq Cr^{-3} (A_1 n^2 + CA_3),$$

where the constant C is independent of n and f .

The estimate (6.3) is a straightforward consequence of formulas (7.18) for G' and (7.14) for ψ_1 , while the case $n = 0$ follows directly from the representation (7.10). \square

Acknowledgments

The research of N. Popivanov and T. Popov was partially supported by the Bulgarian NSF under Grant DCVP 02/1/2009 and by Sofia University Grant 158/2014. The research of A. Tessedal was partially supported under Research Foundation of CUNY Grant 66237-00 44. The authors would like to thank the anonymous referees for the constructive comments and suggestions to improve the quality of the paper.

REFERENCES

- [1] S.A. Aldashev, Correctness of multi-dimensional Darboux problems for the wave equation, *Ukr. Math. J.* 45, No.9 (1993) 1456–1464
- [2] S.A. Aldashev, Spectral Darboux-Protter problems for a class of multi-dimensional hyperbolic equations, *Ukrainian Math. J.* 55, No.1 (2003) 126–135.
- [3] A.K. Aziz, M. Schneider, Frankl-Morawetz problems in R^3 , *SIAM J. Math. Anal.* 10 (1979) 913–921.
- [4] Ar. B. Bazarbekov and Ak. B. Bazarbekov, The Goursat and Darboux problems for the three-dimensional wave equation, *Differ. Equations* 38 (2002) 695–701.
- [5] A.V. Bitsadze, Some classes of partial differential equations, Gordon and Breach Science Publishers, New York, 1988
- [6] E.T. Copson, On the Riemann-Green function, *J. Rath. Mech. Anal.* 1 (1958) 324–348.
- [7] L. Dechevsky, N. Popivanov and T. Popov, Exact asymptotic expansion of singular solutions for (2+1)-D Protter problem, *Abstract and Applied Analysis*, Volume 2012 (2012), Article ID 278542.
- [8] D.E. Edmunds, N.I. Popivanov, A nonlocal regularization of some over-determined boundary value problems I, *SIAM J. Math. Anal.* 29, No.1 (1998) 85–105.
- [9] A. Erdélyi, W. Magnus, F. Oberhettinger, F. Tricomi, Higher transcendental functions, 1, McGraw-Hill, New York, 1953.
- [10] P.R. Garabedian, Partial differential equations with more than two variables in the complex domain, *J. Math. Mech.* 9 (1960) 241–271.

- [11] M.K. Grammatikopoulos, T.D. Hristov, N.I. Popivanov, Singular solutions to Protter's problem for the 3-D wave equation involving lower order terms, *Electron. J. Diff. Eqns.* [online], Vol. 2003 (2003) No.03 pp.1–31. Available from: <http://ejde.math.swt.edu/volumes/2003/03/>
- [12] M. K. Grammatikopoulos, N.I. Popivanov, T. P. Popov, New singular solutions of Protter's problem for the 3 – D wave equation, *Abstract and Applied Analysis* 2004:4 (2004) 315–335.
- [13] L. Hörmander, *The Analysis of Linear Partial Differential Operators III*. Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 1985.
- [14] M. N. Jones, *Spherical Harmonics and Tensors for Classical Field Theory*, Research Studies Press, Letchworth, 1986.
- [15] Jong Duek Jeon, Khe Kan Cher, Ji Hyun Park, Yong Hee Jeon, Jong Bae Choi, Protter's conjugate boundary value problems for the two dimensional wave equation, *J. Korean Math. Soc.* 33 (1996) 857–863.
- [16] Jong Bae Choi, Jong Yeoul Park, On the conjugate Darboux-Protter problems for the two dimensional wave equations in the special case, *J. Korean Math. Soc.* 39, No.5 (2002) 681–692.
- [17] S. Kharibegashvili, On the solvability of a spatial problem of Darboux type for the wave equation, *Georgian Math. J.* 2 (1995) 385–394.
- [18] S. Kharibegashvili, B. Midodashvili, On the solvability of one boundary value problem for one class of semilinear second order hyperbolic systems, *J. Math. Anal. Appl.* 400, No. 2 (2013) 345–362.
- [19] Khe Kan Cher, Nontrivial solutions of some homogeneous boundary value problems for a many-dimentional hyperbolic Euler-Poisson-Darboux equation in an unbounded domain, *Differ. Equations* 34, No.1 (1998) 139–142.
- [20] P. D. Lax, R. S. Phillips, Local boundary conditions for dissipative symmetric linear differential operators, *Comm. Pure Appl. Math.* 13 (1960) 427–455.
- [21] G. Lohöfer, Inequalities for the associated Legendre functions, *Journal of Approximation Theory* 95, No.2 (1998) 178–193.
- [22] D. Lupo, K. Payne and N. Popivanov, Nonexistence of nontrivial solutions for supercritical equations of mixed elliptic-hyperbolic type, *Contributions to nonlinear analysis*, 371–390, "Progress in Non-Linear Differential Equations and Their Applications", 66 Birkhäuser, Basel, 2006.
- [23] C. S. Morawetz, A weak solution for a system of equations of elliptic-hyperbolic type, *Comm. Pure Appl. Math.* 11 (1958) 315–331.
- [24] C. S. Morawetz, Mixed equations and transonic flow, *J. Hyperbolic Differ. Equ.*, Vol. 1, No. 1 (2004) 1–26.
- [25] V. Müller, *Spectral theory of linear operators and spectral systems in Banach algebras*, *Operator Theory: Advances and Applications* 139, Second ed., Birkhäuser Verlag, Basel, 2007.
- [26] N. Popivanov and M. Schneider, The Darboux problems in \mathbf{R}^3 for a class of degenerated hyperbolic equations, *J. Math. Anal. Appl.* 175 (1993) 537–579.
- [27] N. Popivanov and M. Schneider, On M. H. Protter problems for the wave equation in R^3 , *J. Math. Anal. Appl.* 194 (1995) 50–77.
- [28] N. Popivanov, T. Popov, Exact Behavior of Singularities of Protter's Problem for the 3 – D Wave Equation, in: Herzberger, J.(ed), "Inclusion Methods for Nonlinear Problems" with Applications in Engineering, Economics and Physics, *Computing [Suppl.]*, 16 (2002) 213–236.
- [29] N. Popivanov, T. Popov, Singular solutions of Protter's problem for the 3 + 1-D wave equation, *Integral Transforms and Special Functions* 15 No.1, February 2004, 73–91.
- [30] N. Popivanov, T. Popov, R. Scherer, Asymptotic expansions of singular solutions for (3 + 1)-D Protter problems, *J. Math. Anal. Appl.* 331 (2007) 1093–1112.

- [31] N. Popivanov, T. Popov, R. Scherer, Protter-Moravetz multidimensional problems, International Conference on Differential Equations and Dinamical Systems, Proceedings of the Steklov Institute of Mathematics, 278, Issue 1 (2012) 179–198.
- [32] N. Popivanov, T. Popov, A. Tesdall, Semi-Fredholm solvability and asymptotic expansions of singular solutions for Protter problems, AIP Conference Proceedings 1570, 335 (2013) 335–342.
- [33] M.H. Protter, New boundary value problems for the wave equation and equations of mixed type, J. Rat. Mech. Anal. 3 (1954) 435–446.
- [34] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional integrals and derivatives: theory and applications, New York, Gordon and Breach. xxxvi, 1993.
- [35] E.M. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton, N.J., Princeton University Press. XIV, 1970.
- [36] Tong Kwang-Chang, On a boundary-value problem for the wave equation, Science Record, New Series 1, No.1 (1957) 1–3.