

*I pledge that I have neither given nor received
unauthorized assistance during this examination.*

Signature:

- **DON'T PANIC!** If you get stuck, take a deep breath and go on to the next question.
- Unless the problem says otherwise **you must show your work** sufficiently much that it's clear to me how you arrived at your answer.
- No calculators or electronic devices are allowed.
- Unless the problem specifically says, you **do not** need to calculate the final answer. An answer that looks something like

$$\frac{\binom{6}{2}\binom{8}{5}}{\binom{23}{4}} \quad \text{or} \quad \frac{(12)(11)(10) + (11)(10)(9)}{14!}$$

is entirely acceptable.

- You may bring a two-sided sheet of notes on letter-sized paper in your own handwriting.
- There are 10 problems on 8 pages.

Question	Points	Score
1	15	
2	9	
3	6	
4	11	
5	10	
6	10	
7	12	
8	5	
9	1	
10	4	
Total:	83	

Good luck!

[15 points] 1. Suppose we roll a 6-sided die over and over again.

In the following answers, you don't need to compute a final answer, and you may leave an expression like $\binom{13}{6}$ in that form. But you should not leave your answer as an infinite sum.

- (a) What is the expected number of rolls to get a six?

- (b) What is the probability that it takes exactly 5 rolls to get a six?

- (c) What is the probability that it takes 5 or more rolls to get a six?

- (d) What is the probability of getting 2 or fewer sixes in the first 10 rolls?

- (e) What is the probability that the sum of the first 2 rolls is 4 or less?

[9 points] 2. Suppose X and Y have joint density function

$$f(x, y) = \begin{cases} \frac{12}{7}(xy + y^2) & \text{if } 0 \leq x \leq 1 \text{ and } 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find the marginal probability density function of X . **Fully compute all integrals. Be sure to clearly state where the p.d.f. is zero and where it's nonzero.**

(b) Find $P(X < Y)$. **Set up integrals but do not compute them.**

(c) Compute $E(e^{XY^2})$. **Set up integrals but do not compute them.**

- [6 points] 3. Let $X \sim \text{Unif}[0, 2]$. Let $Y = 1/X$. Find the probability density function of Y . If the density is zero outside of some region, be sure to clearly state this.

4. There are 5 types of sandwiches for sale at a shop. For 10 days, you go to the shop for lunch and order one of the 5 sandwiches uniformly at random, independently of previous days.

[4 points]

- (a) What is the probability that you order the falafel sandwich sometime during the 10 days?

[4 points]

- (b) Let X be the number of different sandwiches out of the 5 types you try in the 10 days. Find $E(X)$.

[3 points]

- (c) For $i = 1, \dots, 5$, let X_i be the number of times you order sandwich type i in the 10 days. Are $X_1, \dots, X_5$ independent? Justify your answer.

5. Suppose that $X \sim \text{Poi}(100)$, and recall that $E(X) = 100$ and $\text{Var}(X) = 100$.

[3 points]

(a) Give a bound on $P(X \geq 120)$ using Markov's inequality.

[3 points]

(b) Give a bound on $P(X \geq 120)$ using Chebyshev's inequality.

[4 points]

(c) Recall that if $X_1, \dots, X_{100}$ are independent and $X_i \sim \text{Poi}(1)$, then $X_1 + \dots + X_{100} \sim \text{Poi}(100)$. Using the central limit theorem, estimate

$$P(X_1 + \dots + X_{100} \geq 120),$$

to get another estimate of $P(X \geq 120)$. **Write your answer in terms of $\Phi(x)$, the cumulative distribution function of the standard normal distribution.**

6. Customers arrive at a store. After a customer arrives, the number of minutes until the next customer arrives is random with distribution $\text{Exp}(1)$. The times between customer arrivals are all independent.

A customer arrives. Let X be the number of minutes from now until *two* more customers arrive.

[3 points]

(a) What is $E(X)$?

[3 points]

(b) What is $\text{Var}(X)$?

[4 points]

(c) What is the probability density function of X ?

7. You roll two 4-sided dice. Let U be the minimum and V be the maximum of the two rolls.

[6 points]

- (a) Find the joint probability mass function of U and V . It is easiest to show it as a table, though you can express it any way you want so long as it's clear.

[2 points]

- (b) What is $P(U = V)$?

[2 points]

- (c) What is $P(V = 4 \mid U = 3)$?

[2 points]

- (d) Are U and V independent? Justify your answer.

- [5 points] 8. You have an urn with balls numbered $1, \dots, 10$. You reach in and pull out three balls. You don't consider your three chosen balls to have any order.
- (a) The size of the most natural sample space for the experiment is
☐ 10^3 ☐ $\binom{10}{3}$ ☐ $10 \cdot 9 \cdot 8$ ☐ $3!$
- (b) Let A be the event that your sample includes ball 1 and B the event that your sample contains ball 2. Then A and B are
☐ not disjoint ☐ disjoint
and A and B are
☐ not independent ☐ independent
- (c) Let A be the event that your sample includes balls 1 and 2, and let B be the event that your sample contains ball 3 and 4. Then A and B are
☐ not disjoint ☐ disjoint
and A and B are
☐ not independent ☐ independent
- [1 point] 9. A patient is tested for a rare genetic disease. You model the test probabilistically. Let D be the event that the person truly has the disease. Let N be the event that the person tests negative. Which of the following expressions represents the probability that a person who has tested positive truly has the disease?
☐ $P(D \mid N^c)$ ☐ $P(N^c \mid D)$ ☐ $P(D \cap N^c)$ ☐ $P(D \cup N^c)$
- [4 points] 10. Suppose that X is a continuous random variable with probability density function $f(x)$.
- (a) The range of possible values of X is given by the range of y -values of $f(x)$.
☐ True ☐ False ☐ It depends on $f(x)$
- (b) The range of possible values of X is given by the set of x -values for which $f(x) > 0$.
☐ True ☐ False ☐ It depends on $f(x)$
- (c) For any real number k , it holds that $P(X = k) = 0$.
☐ True ☐ False ☐ It depends on $f(x)$
- (d) For any real numbers a and b with $a < b$, it holds that $P(a < X < b) = P(a \leq X \leq b)$.
☐ True ☐ False ☐ It depends on $f(x)$