

*I pledge that I have neither given nor received
unauthorized assistance during this examination.*

Signature:

- **DON'T PANIC!** If you get stuck, take a deep breath and go on to the next question.
- Unless the problem says otherwise **you must show your work** sufficiently much that it's clear to me how you arrived at your answer.
- No calculators or electronic devices are allowed.
- Unless the problem specifically says, you **do not** need to calculate the final answer. An answer that looks something like

$$\frac{\binom{6}{2}\binom{8}{5}}{\binom{23}{4}} \quad \text{or} \quad \frac{(12)(11)(10) + (11)(10)(9)}{14!}$$

is entirely acceptable.

- You may bring a two-sided sheet of notes on letter-sized paper in your own handwriting.
- There are 9 problems on 11 pages.

Question	Points	Score
1	16	
2	9	
3	6	
4	14	
5	12	
6	6	
7	16	
8	1	
9	7	
Total:	87	

Good luck!

1. A factory makes bolts designed to attach pieces of metal under high pressure. They model their production by assuming that each bolt is defective with probability $1/75$, independent of all other bolts.

[2 points]

- (a) Two pieces of metal are attached with 20 bolts. What is the expected number of defective bolts out of these 20 bolts?

[4 points]

- (b) The two pieces will separate under pressure if 2 or more out of these 20 bolts are defective. What is the probability of this?

[4 points]

- (c) Suppose that the two pieces do not separate under pressure, i.e., that at most 1 of the 20 bolts is defective. Given that this occurs, what is the probability that none of the 20 bolts are defective?

This problem continues onto the next page.

- [3 points] (d) The factory opens in the morning and starts producing bolts. Let X be the number of bolts produced *before* the first defective one. What is $E(X)$? (Note: do not include the defective bolt in your count X . That is, if the first bolt is good and the second is defective, then X is 1 rather than 2.)

- [3 points] (e) For the random variable X from the previous question, find $P(X = 10)$. That is, what is the probability that exactly 10 good bolts are produced before the first defective one?

- [9 points] 2. Let X be the number of hours past noon at which a fire drill takes place. The drill will happen sometime between 1pm and 4pm, but not from 2pm to 3pm, and it is equally likely to occur at any of the permitted times. (That is, X is uniformly distributed on the set $[1, 2] \cup [3, 4]$.)

(a) What is the probability density function of X ?

(b) What is the expectation of X ?

(c) What is the variance of X ?

- [6 points] 3. Suppose that $X_1, \dots, X_{200}$ are i.i.d. with mean 1 and variance 2. Use the normal approximation to estimate $P(X_1 + \dots + X_{200} > 150)$. **Give your answer in terms of $\Phi(x)$, the cumulative distribution function of the standard normal distribution.**

4. A room has bins labeled $1, \dots, 20$. Each of 20 balls is placed into a bin chosen uniformly at random, independently of each other.

[2 points]

- (a) Let X_1 be the number of balls placed into bin #1. What is the **exact** distribution of X_1 ? (Your answer should be a named distribution and you should give its parameter(s).)

[2 points]

- (b) What distribution is appropriate to **approximate** the distribution of X_1 ? (Your answer should be a named distribution and you should give its parameter(s).)

[3 points]

- (c) What is the **exact** probability that bin #1 is empty?

This problem continues onto the next page.

[3 points] (d) What is the expected number of empty bins?

[4 points] (e) What is the probability that every bin contains exactly one ball?

[12 points] 5. Suppose that X and Y have joint probability density function

$$f(x, y) = \begin{cases} \frac{3}{2}(x^2 + y^2) & \text{if } 0 \leq x \leq 1 \text{ and } 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) Compute the marginal probability density function of X . **Fully compute all integrals in this answer.**

- (b) Are X and Y independent? Justify your answer.

This problem continues onto the next page.

- (c) Set up an integral to find $P(Y > X)$. **Your answer should be a parametrized, double integral, but don't compute it.**
- (d) Find $P\left(Y > \frac{1}{2} \mid Y > X\right)$. **Your answer should be an expression of parametrized double integrals, and you should not compute them.**

[6 points] 6. Suppose that $X_1, X_2, \dots$ are independent and identically distributed, with $E(X_i) = 6$ and $\text{Var}(X_i) = 2$.

(a) Use Chebyshev's inequality to bound $P(|X_1 - 6| > 4)$.

(b) Compute the mean and variance of $X_1 + \dots + X_{10}$.

7. Flip a fair coin four times. Let X be the number of tails in flips 1, 2, and 3. Let Y be the number of tails in flips 3 and 4.

[4 points]

- (a) Give the **marginal** distributions of X and Y . (Your answers should be named distributions and you should give their parameter(s).)

[4 points]

- (b) Compute $P(X = 2, Y = 1)$.

[4 points]

- (c) Compute $P(X = 2 \mid Y = 1)$.

[3 points]

- (d) Are X and Y independent or not? Justify your answer rigorously, with more than just intuition.

[1 point]

- (e) Which of the following is correct? (You do not need to justify your answer. You do not need to do any calculations to answer this question, if you use your intuition correctly.)

☐ $\text{Cov}(X, Y) > 0$

☐ $\text{Cov}(X, Y) < 0$

☐ $\text{Cov}(X, Y) = 0$

- [1 point] 8. A patient is tested for a rare genetic disease. You model the test probabilistically. Let D be the event that the person truly has the disease. Let N be the event that the person tests negative. Which of the following expressions represents the probability that a person who has tested positive truly has the disease?

☐ $P(D \cap N^c)$ ☐ $P(D \cup N^c)$ ☐ $P(D \mid N^c)$ ☐ $P(N^c \mid D)$

- [7 points] 9. Say whether the following statements are true or false.

(a) If events A and B are disjoint, then the event $A \cap B$ never occurs.

☐ True ☐ False

(b) If events A and B are disjoint, then $P(A \cup B) = P(A) + P(B)$.

☐ True ☐ False

(c) If events A and B are independent, then $P(A \cap B) = P(A) + P(B)$.

☐ True ☐ False

(d) It makes sense to talk about events A and B being disjoint, and it also makes sense to talk about random variables X and Y being disjoint.

☐ True ☐ False

(e) It makes sense to talk about events A and B being independent, and it also makes sense to talk about random variables X and Y being independent.

☐ True ☐ False

(f) If random variables X and Y are independent, then their distributions are always discrete.

☐ True ☐ False

(g) The expectation of any random variable is a positive number.

☐ True ☐ False